

## **UNDERSTORY VEGETATION INFLUENCES INSECT DIVERSITY IN RUBBER PLANTATIONS OF KANYAKUMARI, INDIA**

### **ABSTRACT:**

Insects play a crucial role in maintaining ecological balance, and their decline has harmful effects on various organisms. Globally, the establishment of rubber plantations have been associated with a reduction in insect diversity. However, the impact of monoculture rubber plantations on biodiversity in India, particularly in Kanyakumari, is not well understood and has received little research attention. This study aimed to assess the status of insect diversity in rubber plantations in this region. Kanyakumari, Tamil Nadu, with its extensive monoculture rubber plantations, provides an ideal setting for this investigation. Three adjacent rubber plantations with varying topography and understory vegetation were selected for the study. Results indicated that insect diversity was significantly higher in Plantation-1 compared to the other two plantations. Canopy and understory vegetation were identified as key factors influencing insect diversity. Plantations with dense understory vegetation consisting of diverse native plant species exhibited greater insect richness. The predominant insect orders across all plantations were Hymenoptera and Diptera. However, the impact of rubber plantations on individual insect species varied based on their habitat preferences. Seasonal fluctuations in diversity were particularly noticeable during the monsoon season. Further comparative studies are needed to understand the broader implications of rubber plantations on insect diversity across the district.

**KEYWORDS:** Insect diversity, Monoculture rubber plantations, Kanyakumari, Seasonal fluctuations

### **Introduction:**

Insects are integral to ecosystem functions such as pollination, pest control, and nutrient cycling, yet their populations are declining globally due to habitat loss and agricultural expansion [1]. The objectives of our study were threefold: to identify the diversity of insects in these plantations, to observe seasonal variations, and to determine the effects of physicochemical parameters of soil and water on insect diversity.

Kappukadu, with its extensive rubber plantations, provides a unique opportunity to study these dynamics in a region where rubber cultivation plays a significant economic role. This research is particularly relevant in the context of the United Nations Sustainable Development Goals (SDGs), particularly “Goal 15: Life on Land”, which emphasizes the need to protect, restore, and promote sustainable use of terrestrial ecosystems. By understanding insect diversity and its determinants, we contribute to the broader goals of conserving biodiversity and promoting sustainable agricultural practices.

Studying these sites is crucial for several reasons. First, it helps in assessing the impact of rubber monoculture on local biodiversity, providing insights into how such practices can be managed to mitigate negative ecological effects. Second, the findings can inform better

**Comment [s1]:** Consider adding a brief explanation of why understory vegetation is particularly important for insect diversity, as this is mentioned in your title but not clearly in the introduction.

**Comment [s2]:** Include some specific numerical data to support your main findings. For example, you could mention the number of insect species or orders found, or the difference in diversity indices between plantations.

**Comment [s3]:** Change "harmful effects" to "detrimental effects" for more formal scientific language.

**Comment [s4]:** Briefly mention the methods used for data collection and analysis. This could be done in one concise sentence.

agricultural practices that balance economic benefits with ecological sustainability. Third, understanding the seasonal variations and environmental factors affecting insect populations can lead to more effective conservation strategies, ensuring that these vital organisms continue to support ecosystem services critical for human well-being and agricultural productivity [2,3]

Studies on insect populations in rubber plantations have been conducted globally, particularly in Southeast Asia. These studies are significant because they provide insights into the impact of monoculture plantations on biodiversity. For instance, research in Thailand, Malaysia, and Indonesia has highlighted how rubber plantations affect insect diversity, comparing these ecosystems with natural forests. These studies have shown that rubber monocultures typically support fewer species and lower insect abundance compared to more diverse habitats [4 – 10].

This underscores the ecological costs of expanding rubber cultivation. The findings have prompted calls for more sustainable agricultural practices, such as integrating agroforestry and maintaining patches of natural vegetation within plantations to support biodiversity. These efforts align with the principles of sustainable development and conservation.

However, comprehensive studies on the impact of rubber plantations on insect biodiversity are less common in India, despite the country's significant rubber production [11 – 12]. This gap may be due to limited funding, lack of awareness, or prioritization of economic benefits over ecological considerations. Nonetheless, the growing recognition of biodiversity's role in ecosystem services and agricultural productivity is driving more research and policy changes aimed at balancing agricultural development with ecological sustainability.

## **2. METHODOLOGY:**

The selected rubber plantations were located in Kappukadu village, Vilavancode taluk, Kanyakumari District, Tamil Nadu, India. These sites are situated between 8°17'14" N and 8°17'9" N latitude and 77°11'55" E and 77°11'57" E longitude. Three plantations were chosen for the study, with areas of 3 hectares, 1.5 hectares, and 2 hectares, respectively.

Plantation 1 (Fig. 1) featured dense understory vegetation and a substantial litter layer. Within this plantation, there was a site with newly planted rubber saplings intercropped with plantain. Plantation 2 had sparse shrubs and lacked grass cover (Fig. 1), with a thin litter layer and natural water springs. Plantation 3 (Fig. 1) was located near a residential area, characterized by a minimal litter layer and scattered shrubs.



**Fig 1.**

**Comment [s5]:** Include caption

## 2.1 Sampling method:

Data were collected over seven months, from July 2021 to January 2022, with weekly visits to each plantation between 6 am and 8 am. Foliage-dwelling and fast-moving insects were observed using the direct observation method. For fast-flying insects, such as dragonflies, random sampling and counting were conducted. Quadrats of 2 m  $\times$  2 m were placed in various locations within each plantation to record the number of slow-moving insects. Sticky traps were hung in different spots of each plantation in the evening and checked the next morning, with the trapped insects collected. Pitfall traps, made using cups and bottles, were buried at different locations in the plantations; water was used instead of killing agents to prevent insects from escaping. Containers filled with water were placed under light bulbs in various locations to trap nocturnal insects. Insect samples were also collected by digging soil and leaf litter and handpicking. The collected insects were examined under a microscope and photographed [13 – 15].

## 2.2 Identification:

The insects were identified using Insect identification manual by ZSI and online insect identification app (iNaturalist) [16,17].

## 2.3 Data analysis:

**Comment [s6]:** Provide more details on the statistical methods used, including any software packages.

The data was subjected to Simpson [18] and Shannon-Weiner index calculation [19] to measure species diversity.

## 2.4 Seasonal variation:

The seasonal data (temperature and rainfall) of the study location for 7 months (July 2021 to January 2022) was collected from [Climate-Data.org website](https://climate-data.org) [20]. Correlation between Insect diversity and seasonal data was found using Pearson's correlation coefficient.

**Comment [s7]:** Italicize websites. Check author guidelines for clarity

**Formatted:** Font: Italic

## 3. RESULTS AND DISCUSSION:

Insects from 13 different orders were recorded (Fig. 2). A total of 87 insect species, spanning 50 families, were observed within these orders (Fig. 2). The order Hymenoptera had the highest number of identified insects, accounting for 29.7% of the total. In contrast, the orders Dermaptera, Mecoptera, and Neuroptera had the lowest number of insects collected.




Figure.2 Diversity of Insects in the rubber plantations

The highest species richness was observed in Lepidopterans. A total of 21 Lepidopteran species from 9 families were recorded, with most sightings occurring in Plantation 1 (Table 1). Hymenoptera was the most abundant insect order observed, with a total of 942 individuals recorded during the study period. The highest number of insects was recorded in August (Table 2).

Plantation 1 exhibited higher diversity, with a Shannon diversity index of 1.81 and a Simpson's index of 0.79, attributed to its dense understory vegetation. This plantation had a greater number of Hymenopterans and Dipterans, with ants being particularly abundant. Both arboreal and terrestrial nesting ant species were recorded. Arboreal ants, such as weaver ants



(*Oecophylla smaragdina*), built their nests on rubber trees in the plantation. Weaver ants (Fig. 3) were found across all three plantations, demonstrating resilience to disturbances in these environments. The closed canopy and thick litter layer in Plantation 1 supported various ant species, with groups of weaver ants observed building nests and foraging (Fig. 3). Among the Diptera, mosquitoes were notably abundant, especially during the monsoon season due to the availability of breeding water. Mosquito larvae were frequently found in rubber latex collecting cups and plastic bottles filled with rainwater in the plantation (Fig. 3).




Figure.3 Mosquito larva in latex collection cup and *Oecophylla smaragdina* on Rubber tree



Figure.4 *Luprops tristis* and *Gesonula punctifrons* on Basket grass



Figure.5 Termite mounds, basket grass and *Leptocoris oratoria*

Plantation 2 had minimal understory vegetation and a thin litter layer. Insects from 9 orders were collected from this plantation, which showed lower diversity with a Shannon diversity index of 1.53 and a Simpson's index of 0.71. Litter-dwelling beetles and ants were absent,

and only two coleopteran species were found during the study period. Plantation 2 features two natural springs, which provide a consistent water source except during the dry summer months. The presence of water supported Odonates (Fig. 8). Weaver ants were also abundant in this plantation, and Hymenopterans were the most frequently collected insects. In contrast, Coleoptera numbers were the lowest, likely due to the lack of understory vegetation, which may have contributed to the reduced richness and diversity of coleopterans. No individuals from the orders Mecoptera, Mantodea, Neuroptera, or Dermaptera were recorded.

Plantation 3 was the second most diverse of the three plantations, with a Shannon diversity index of 1.73 and a Simpson's index of 0.78. Insects from 8 orders were observed. Although the total number of individuals was lower than in Plantation 2, the distribution of species was more even. Dipteran insects were the most abundant, followed by Hymenoptera, while Blattodea had the fewest individuals. No insects from the orders Dermaptera, Mecoptera, Mantodea, or Neuroptera were recorded. Mosquitoes were particularly dominant in this plantation, likely due to the presence of nearby residents and associated anthropogenic activities, which may have contributed to their increased diversity and abundance. The leaf litter was sparse, and moderate understory vegetation was present.

Table.1 Insect species recorded from Monoculture Rubber plantations in Kanyakumari

S <sub>2</sub> No <sub>2</sub>	Order	Family	Organism	Observed in
1.	Blattodea	Blattidae	<i>Periplaneta fuliginosa</i>	Plantation 1, 2 and 3
		Ectobiidae	<i>Blattella asahinai</i>	Plantation 1
			<i>Supella longipalpa</i>	Plantation 3
			<i>Ectobius vittiventris</i>	Plantation 1 and 3
2.	Coleoptera	Coccinellidae	<i>Delphastus pusillus</i>	Plantation 1
			<i>Coccinella septempunctata</i>	Plantation 1 and 3
		Chrysomelidae	<i>Aulacophora foveicollis</i>	Plantation 1, 2 and 3
			<i>Luperus flavipes</i>	Plantation 1
			<i>Mordella marginata</i>	Plantation 1
			<i>Heteronychus arator</i>	Plantation 1
			<i>Hydnobius punctatus</i>	Plantation 1
			<i>Litargus connexus</i>	Plantation 1
			<i>Luprops tristis</i>	Plantation 1, 2 and 3
			<i>Luciola lateralis</i>	Plantation 1, 2 and 3

				3
			<i>Hydaticus aruspex</i>	Plantation 2
3.	Dermaptera	Forficulidae	<i>Forficula auricularia</i>	Plantation 1
4.	Diptera	Bombyliidae	<i>Poecilanthrax apache</i>	Plantation 1
		Micropezidae	<i>Rainiera antennaepes</i>	Plantation 1, 2 and 3
		Chironomidae	<i>Chironomus plumosus</i>	Plantation 1, 2 and 3
		Culicidae	<i>Aedes aegypti</i>	Plantation 1, 2 and 3
		Muscidae	<i>Musca domestica</i>	Plantation 1, 2 and 3
		Calliphoridae	<i>Lucilia sericata</i>	Plantation 1
		Tephritidae	<i>Tephritis conura</i>	Plantation 1
			<i>Anomoia purmunda</i>	Plantation 1
		Tipulidae	<i>Dolichocheza walleyi</i>	Plantation 1
5.	Hemiptera	Alydidae	<i>Leptocoris oratoria</i>	Plantation 1
		Lygaeidae	<i>Oncopeltus fasciatus</i>	Plantation 1 and 3
		Miridae	<i>Lygocoris pabulinus</i>	Plantation 1, 2 and 3
		Cercopidae	<i>Cercopis saguinolenta</i>	Plantation 1 and 2
		Cicadellidae	<i>Aphrodes bicintus</i>	Plantation 1, 2 and 3
6.	Hymenoptera	Sphecidae	<i>Chalybion californicum</i>	Plantation 1
		Apidae	<i>Apis mellifera</i>	Plantation 1 and 3
			<i>Amegilla cingulata</i>	Plantation 1
		Halictidae	<i>Halictus farinosus</i>	Plantation 1 and 3
			<i>Augochlora pura</i>	Plantation 1, 2 and 3
		Formicidae	<i>Oecophylla smaragdina</i>	Plantation 1, 2 and 3
			<i>Odontomachus bauri</i>	Plantation 1

			<i>Pogonomyrmex bicolor</i>	Plantation 1
			<i>Camponotus radiates</i>	Plantation 1, 2 and 3
			<i>Tetraponera allaborans</i>	Plantation 1
			<i>Anoplolepis gracileps</i>	Plantation 1
			<i>Monomorium pharaonic</i>	Plantation 1, 2 and 3
			<i>Paratrechina longicornis</i>	Plantation 1, 2 and 3
		Pompilidae	<i>Auplopus carbonarius</i>	Plantation 1
7.	Isoptera	Rhinotermitidae	<i>Coptotermes formosanus</i>	Plantation 1, 2 and 3
8.	Lepidoptera	Uraniidae	<i>Micronia aculeate</i>	Plantation 1
		Crambidae	<i>Achyra ranatlis</i>	Plantation 1
			<i>Nausinoe geometralis</i>	Plantation 1
			<i>Patania ruralis</i>	Plantation 1
			<i>Sphragedius similis</i>	Plantation 1
		Erebidae	<i>Syntomoides imaon</i>	Plantation 1
		Pterophoridae	<i>Hellinsia pectodactylus</i>	Plantation 1
		Nymphalidae	<i>Ypthima huebneri</i>	Plantation 1 and 3
			<i>Junio lemonias</i>	Plantation 1
			<i>Mycalesis perseus</i>	Plantation 2 and 3
		Papilionidae	<i>Troides minos</i>	Plantation 1
			<i>Battus polydamus</i>	Plantation 1
			<i>Papilio polytes</i>	Plantation 1
		Lycaenidae	<i>Talicauda nyseus</i>	Plantation 1
			<i>Zizula hylax</i>	Plantation 1
			<i>Euchrysops cnejus</i>	Plantation 1
			<i>Jamides celeno</i>	Plantation 1
			<i>Castalius rosimon</i>	Plantation 1



		Hesperiidae	<i>Lambrix salsa</i>	Plantation 1
			<i>Arnetta vindhiana</i>	Plantation 1
		Pieridae	<i>Eurema blanda</i>	Plantation 1, 2 and 3
9.	Mantodea	Liturgusidae	<i>Liturgusa maya</i>	Plantation 1
		Mantidae	<i>Hierodula patellifera</i>	Plantation 1
			<i>Ameles decolor</i>	Plantation 1
10.	Mecoptera	Panorpidae	<i>Panorpa nuptialis</i>	Plantation 1
11.	Neuroptera	Myrmeleontidae	<i>Distoleon tetragamicus</i>	Plantation 1
12.	Odonata	Coenagrionidae	<i>Ceragrion cerinorubellum</i>	Plantation 1
			<i>Pesudagrion microcephalum</i>	Plantation 1
		Platycnemididae	<i>Copera marginipes</i>	Plantation 1
		Libellulidae	<i>Rhyothemis variegata</i>	Plantation 1
			<i>Neurothemis tullia</i>	Plantation 1 and 2
			<i>Trithemis aurora</i>	Plantation 1, 2 and 3
			<i>Tholymis tillarga</i>	Plantation 1 and 2
			<i>Orthetrum glaucum</i>	Plantation 2
			<i>Orthetrum chrysis</i>	Plantation 2
			<i>Diplacodes trivalis</i>	Plantation 1
13.	Orthoptera	Acrididae	<i>Gesonula punctifrons</i>	Plantation 1
		Gryllidae	<i>Xenogryllus marmortus</i>	Plantation 1
			<i>Xenogryllus sp</i>	Plantation 1
			<i>Telogyllus emma</i>	Plantation 1
		Tetrigidae	<i>Tetrix tenuicornis</i>	Plantation 1
			<i>Paratettix curtispennis</i>	Plantation 2 and 3

MONTHS	JULY			AUGUST			SEPTEMBER			OCTOBER			NOVEMBER			DECEMBER			JANUARY		
INSECTS	P-1	P-2	P-3	P-1	P-2	P-3	P-1	P-2	P-3	P-1	P-2	P-3	P-1	P-2	P-3	P-1	P-2	P-3	P-1	P-2	P-3
Blattodea	5	7	3	12	-	-	11	-	1	1	1	-	1	-	-	2	4	-	-	3	1
Coleoptera	21	-	5	21	-	7	35	-	11	22	2	9	29	-	4	11	-	2	6	-	-
Dermaptera	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	2	-	-
Diptera	86	32	10	123	30	8	103	45	29	56	31	18	144	57	25	69	10	14	17	15	11
Hemiptera	7	-	-	2	-	-	6	1	-	1	2	5	58	4	9	29	25	-	2	-	-
Hymenoptera	102	37	9	170	78	13	173	44	7	48	34	9	66	33	3	7	10	4	52	26	17
Isoptera	-	10	5	152	-	-	18	-	8	-	-	-	-	-	-	-	4	-	25	20	10
Lepidoptera	24	4	4	48	3	2	56	2	2	14	2	1	20	1	NO	6	2	2	24	5	9
Mantodea	-	-	-	1	-	-	1	-	-	-	-	-	0	-	-	-	-	-	1	-	-
Mecoptera	-	-	-	-	-	-	-	-	-	-	-	-	2	-	-	-	-	-	-	-	-
Neuroptera	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	-	-
Odonata	6	1	-	8	2	-	7	2	-	2	3	-	1	4	-	-	4	-	2	3	-
Orthoptera	16	2	-	14	5	-	15	7	3	3	9	7	17	14	5	11	6	6	5	2	-
Total	267	93	36	551	118	30	425	101	61	147	84	49	338	113	46	136	65	28	138	74	48

Table.2 Number of Insects recorded in each insect order

**Comment [s8]:** Table caption should be placed above the table



Figure.6 *Papilio polytes* and *Jamides celeno*



Figure. 7 *Teleogryllus emma*




Figure.8 Odonates from Plantation 2, *Neurothemis tullia* and *Tholymis tillarga*

### 3.2 SEASONAL VARIATION:

**Comment [s9]:** Follow lower case for subheadings and mention 3.1 sub heading



**Figure.9** Seasonal variation in vegetation, Canopy and litter during January in Plantation -I



The understory vegetation and canopy density varied throughout the year. From June to November, the canopy was dense, while both the understory and canopy were less dense in December and January, though the litter layer increased in thickness (Fig. 9). The highest temperature was recorded in September, and the lowest in January 2022. Rainfall peaked in

November 2021, with 183 mm recorded, whereas January 2022 saw the lowest rainfall (Fig. 10). Temperature positively influenced insect diversity in Plantations 1 and 2, but Plantation 3 exhibited a negative correlation with temperature (Fig. 11, Table 3). This is supported by the study conducted on the litter insects and their relationship with rainfall in Western Ghats which states, seasonal variation was not seen on the whole. But, orthopteran numbers increased during monsoon [21].




Figure.12 Correlation between Insect diversity and Temperature

Table.3 Pearson's Correlation coefficient

PLANTATIONS	TEMPERATURE (in C)	RAINFALL (in mm)
1	0.56481	-0.12557
2	0.550524	0.226639
3	-0.01947	0.174588

**Comment [s10]:** Try to include tables in editable format for clear appearance, instead of picture



Rainfall positively affected insect diversity in Plantations 2 and 3, whereas Plantation 1 showed a negative correlation with rainfall (Fig. 12, Table 3). However, the diversity of some insect groups remained relatively stable despite seasonal changes. This finding aligns with a study on litter insects in the Western Ghats, which reported minimal overall seasonal variation but noted an increase in orthopteran numbers during the monsoon.




Figure.10 Temperature and Rainfall Data




Figure.11 Correlation between Insect diversity and Temperature

The study's data and analyses reveal insights into insect diversity across three plantations, highlighting seasonal variations and the impact of temperature and rainfall. Temperature remains relatively stable from July to January, while rainfall peaks in October and November.



The correlation between temperature and insect numbers shows a moderate positive relationship in Plantations 1 and 2, with coefficients of 0.56481 and 0.550524, respectively, indicating that higher temperatures are associated with more insects. Plantation 3 shows no significant correlation (-0.01947). Conversely, rainfall exhibits weaker and more variable correlations with insect numbers. In Plantation 1, there is a weak negative correlation (-0.12557), while Plantations 2 and 3 show weak positive correlations (0.226639 and 0.174588, respectively). These findings suggest that temperature is a more consistent factor influencing insect diversity, particularly in Plantations 1 and 2, whereas the impact of rainfall is less clear and more variable across the plantations.

#### 4. CONCLUSION:

Of the 81 insects recorded, 47 were found exclusively in Plantation 1, highlighting that understory vegetation with natural flora can significantly enhance insect biodiversity, even in monoculture rubber plantations. Specific plants attracted various insects; for instance, *Clerodendrum infortunatum* was prevalent in Plantation 1, attracting species such as the *Troides minos* butterfly, *Tephritis conura* and *Anomoia purmunda* fruitflies, and *Oecophylla smaragdina* (weaver ants) which were observed collecting nectar from its flowers. Similarly, plants like basket grass supported several insects, including *Tetraponera allaborans* (slender ants) and *Leptocoris oratoria* (rice ear bug). This indicates that native plants can enhance insect diversity within rubber plantations.

Maintaining healthy understory vegetation with a mix of native shrubs and grasses positively impacts insect diversity. Effective understory management practices, such as limiting the collection of twigs and litter and avoiding excessive weeding of native grasses, can support ground-dwelling and leaf-eating insects. Currently, many local rubber plantation workers and owners are unaware of the impact of their practices on biodiversity. Educating them about the benefits of these practices could lead to improved overall insect biodiversity.

#### REFERENCES:

1. Footitt, R. G., & Adler, P. H. (Eds.). (2009). Insect biodiversity: science and society. John Wiley & Sons.
2. Samways, M. J. (2015). Future-proofing insect diversity. *Current opinion in insect science*, 12, 71-78.
3. Jankielsohn, A. (2018). The importance of insects in agricultural ecosystems. *Advances in Entomology*, 6(2), 62-73.
4. Aratrakorn S, Thunhikorn S, Donald PF (2006) Changes in bird communities following conversion of lowland forest to oil palm and rubber plantations in southern Thailand. *Bird Conserv Int*. 16: 71-82
5. Phommexay, P., Satasook, C., Bates, P., Pearch, M., & Bumrungsri, S. (2011). The impact of rubber plantations on the diversity and activity of understorey insectivorous bats in southern Thailand. *Biodiversity and Conservation*, 20(7), 1441-1456.

**Comment [s11]:** Consider adding recommendations for future research or management practices based on your findings.

**Comment [s12]:** Follow Author guidelines for all the references. Write either full names of journal or in abbreviation form, as per the author guidelines

6. Pereira, H. M., Navarro, L. M., & Martins, I. S. (2012). Global biodiversity change: the bad, the good, and the unknown. *Annual Review of Environment and Resources*, 37, 25-50.
7. Meng, LZ., Martin, K., Weigel, A. *et al.* Impact of rubber plantation on carabid beetle communities and species distribution in a changing tropical landscape (southern Yunnan, China). *J Insect Conserv* **16**, 423–432 (2012).  
<https://doi.org/10.1007/s10841-011-9428-1>
8. Liu, C. L. C., Kuchma, O., & Krutovsky, K. V. (2018). Mixed-species versus monocultures in plantation forestry: Development, benefits, ecosystem services and perspectives for the future. *Global Ecology and Conservation*, 15, e00419
9. Lee RH, Wang CL-W, Guénard B. The ecological implications of rubber-based agroforestry: Insect conservation and invasion control. *J Appl Ecol*. 2020; 57: 1605–1618. <https://doi.org/10.1111/1365-2664.13642>
10. Singh Ashutosh Kumar, Wenjie Liu, Sissou Zakari, Junen Wu, Bin Yang, Xiao Jin Jiang, Xiai Zhu, Xin Zou, Wanjun Zhang, Chunfeng Chen, Rishikesh Singh, A run Jyoti Nath (2021) A global review of rubber plantations: Impacts on ecosystem functions, mitigations, future directions, and policies for sustainable cultivation. *Science of The Total Environment* Vol. 796 pg. 148948 DOI - <https://doi.org/10.1016/j.scitotenv.2021.148948>
11. Larjavaara, M. (2008). A review on benefits and disadvantages of tree diversity. *The Open Forest Science Journal*, 1(1).
12. Meti, S., Pradeep, B., Jacob, J., Meerabai, M. and Jessy, M.D. (2014). Spatio-temporal analysis of rubber area and its association with soil and topography in Kanyakumari district. *Rubber Science*, 27(2):182-192.
13. Nyffeler, M. (1982). Field studies on the ecological role of the spiders as insect predators in agroecosystems (abandoned grassland, meadows, and cereal fields) (Doctoral dissertation, ETH Zurich).
14. Wiebe, K. L., & Gow, E. A. (2013). Choice of foraging habitat by Northern Flickers reflects changes in availability of their ant prey linked to ambient temperature. *Ecoscience*, 20(2), 122-130.
15. Kyerematen, R., Acquah-Lamptey, D., Owusu, E. H., Anderson, R. S., & Ntiama Baidoo, Y. (2014). Insect diversity of the muni-pomadze Ramsar site: An important site for biodiversity conservation in Ghana. *Journal of Insects*, 2014.
16. iNaturalist. (n.d.). Retrieved from <https://www.inaturalist.org/>
17. ZSI resources from: <https://faunaofindia.nic.in/>
18. SIMPSON, E. Measurement of Diversity. *Nature* 163, 688 (1949). <https://doi.org/10.1038/163688a0>
19. Shannon, C.E. and Weaver, H.J. (1949). *The Mathematical theory of communication* University of Illinois press, Urbana
20. <https://en.climate-data.org/asia/india/tamil-nadu/kanniyakumari-34132/>
21. Anu, A., Sabu, T. K., & P.J. V. (2009). Seasonality of litter insects and relationship with rainfall in a wet evergreen forest in south Western Ghats. *Journal of Insect Science*, 9(1), 46.