

Diversity and Relative Abundance of Mosquito (Diptera: Culicidae) within human habitat in the district of Thane, Maharashtra, India

Abstract

Mosquito's form an integral part of our ecosystem. They act as **vector** in transmitting diseases and weakening the human immune system. A large amount of the country's economy is expended in controlling the growth and spread of mosquitoes and the associated diseases. In the current study dead mosquitoes from different areas of Mira Bhayander Thane district were collected and identified using standard identification methods. The study was focused on the three genera *Aedes*, *Anopheles* and *Culex* as they are most commonly found in Mumbai and Thane region and are responsible for diseases like dengue, malaria, filariasis, etc which are currently on the rise. A total of 1123 dead mosquitoes were collected and it was observed that the **population** of *Culex* was **highest** and *Anopheles* was the least. Relative abundance of species was also calculated which concluded that the *Aedes* and *Culex* mosquitoes were the dominant **genera** whereas the *Anopheles* was the subdominant **genera**. The present study will aid in limiting the spread of diseases by controlling the growth of *Culex* and other mosquitoes.

Keywords

Aedes, *Anopheles*, *Culex*, Mira Bhayander

1. Introduction

Mosquitoes have bedeviled humans for centuries, spreading diseases and **caused death to millions of people**. Mosquitoes are chief **vector** that are accountable for causing diseases moreover their diversity and distribution determine the potency of disease transmission and also the ecological status of surrounding environment [1]. **They act as pathogen vector that caused disease in humans and livestock similarly**. However, mosquitoes also have an importance ecologically, not only are they efficient pollinators but the larva of mosquitoes serves as a source of food for variety of fishes and birds that feed on very small aquatic animals [2]. There are 112 genera of mosquitoes worldwide out of which only few bites and play a role in causing diseases in humans [3]. According to WHO, vector borne diseases are 17% of all the infectious diseases that are caused globally out of which approximately **7 lakh** every year results in death [4]. *Aedes*, *Anopheles* and *Culex* genera are the most common and potential vectors for parasites that can cause various diseases like dengue, malaria, encephalitis, lymphatic filariasis

and West Nile virus [5,6]. Mosquitoes not only play a role in transmitting various pathogens but also aid in suppressing the human immune system. As soon as the mosquito saliva enters to the human blood, it suppresses the production of cytokines in human peripheral blood mononuclear cells, thus helping the pathogen make us infected [7]. It also reduces IP10 which attracts macrophages, monocytes and dendritic cells[8]. Within the period of 7 days it also reduces neutrophils and myeloid cells that could suppress the growth of pathogens in the blood. Thus, adversely affecting the human immune system [9,10]. During Covid 19, people were strengthening their immune system in order to not get infected by the virus [11, 33-34]. This could be the reason that the cases of mosquito borne diseases reduced during the late 2020's. There are several records of studies done to understand the diversity and abundance of mosquitoes and mosquito borne diseases worldwide as well as in India, but there is barely any study that represents the diversity and abundance of mosquitoes in Maharashtra. The present work was an attempt to study the prevalence of three commonly found genera *Aedes*, *Culex* and *Anopheles* and the diseases concerned with them. According to 2011 census, Mira Bhayander has a population of approximately 8 lakh people [12]. As per the severity of this disease, the cases were expected to be quite high comparatively, with 877 cases of filariasis within a span of 6 months i.e. January to June 2020 [13]. According to WHO, at least 36 million people live with the manifestation of this disease [4]. Hydrocele and lymphoedema are symptoms of filariasis [4] however, due to lack of awareness, the people are not aware of the gravity of disease and detailed symptoms or causes. The disease is not deadly but causes unnecessary expense to treat the symptoms that are increased because of negligence. The present study will help to understand the intensity of infection in areas of Mira Bhayander and to adapt methods for the spread of disease.

2. Materials and Methods

2.1. Sampling site

The study was conducted in Mira Bhayander which is located in Thane district, Maharashtra, India. It can be located using GPS co-ordinates 19° 17' 42.8388'' N and 72°51' 15.8148'' E. The study was carried out in four different locations i.e. Rai, Navgarh Road, MBMC Dumping site and Uttan of Mira Bhayander for collection of samples that were close to human habitat and also to places that could be the possible mosquito breeding sites.

2.2. Sampling method and identification

The collection of samples of mosquito started in the month of July 2022 and ended in December 2022, it included monsoon and winter seasons. Due to fluctuations in the temperature during the month of October, it was the most efficient time to study about the impact of temperature

on the diversity of mosquitoes. The mosquito samples were collected by human landing method and mostly by light trap method using UV LED mosquito trap machine between dawn and dusk i.e. between 8:00 pm to 7:00 am placed at restaurants and stalls. The trapped mosquitoes were collected in a vial and then taken to the laboratory of the Zoology Department of Bhavan's College (Autonomous), Andheri west, Mumbai for identification. The samples were then stored in formalin until it was further identified and sorted. The identification was done with the help of compound microscope and dissecting microscope based on standard taxonomic keys- A catalogue of Indian Mosquitoes [9]. The mosquitoes were first identified as male and female and then on the basis of their genera. The current work did not involve any breeding or collection of live mosquitoes, only dead mosquitoes were collected for the study.

2.3.Data Analysis

The relative abundance (RA) of mosquitoes [15] was calculated by using the formula given by Weiher and Keddy.

$$RA = \frac{l}{L} \times 100$$

• Where l is equal to the number of specimens collected of particular species and L is equal to the total number of specimens collected. Based on the values of RA, the specimens are considered as satellite (RA < 1%); sub-dominant (RA < 5%) and dominant species (RA > 5%).

Shannon diversity index (H') and Simpson's diversity index (D) was calculated in order to find out diversity of mosquito in the district using Microsoft Excel Worksheet (Version 2016) [15].

Shannon diversity index (H);

$$H = - \sum_{i=1}^s p_i \ln p_i$$

Simpson diversity index (D);

$$D = \frac{N(N-1)}{\sum n(n-1)}$$

3. Results

A total of 1123 mosquitoes were collected out of which 661 were identified as *Culex* (58.86%), 328 *Aedes* (29.2%), 23 *Anopheles* (2.05%) and 111 belong to other genera which were not considered for the study (9.88%). According to the theory of Weiher and Keddy, relative abundance was calculated as per which *Anopheles* was considered to be subdominant

species with RA= 2.04, *Culex* and *Aedes* were considered to be dominant species with RA equals 58.86 and 29.20 respectively. Thus within the duration of 6 months it was found that the genus *Culex* was the dominating species as compared to other disease causing mosquitoes.

As per the Simpson's index and the Shannon- Wiener index; *Anopheles* was found to be moderately diverse whereas *Culex* and *Aedes* was found to be highly diverse. The correlation between the two index

Table 1. Diversity and Relative Abundance of Mosquitoes in Mira Bhayander of Thane District

| | Site 1 | Site 2 | Site 3 | Site 4 | Total | RA (%) | Status |
|------------------|--------|--------|--------|--------|-------------|--------|--------------|
| <i>Anopheles</i> | 3 | 6 | 2 | 12 | 23 | 2.04 | Sub dominant |
| <i>Aedes</i> | 88 | 33 | 193 | 347 | 661 | 58.8 | Dominant |
| <i>Culex</i> | 63 | 7 | 9 | 249 | 328 | 29.2 | Dominant |
| Others | 25 | 11 | 5 | 70 | 111 | 9.88 | Dominant |
| Total | | | | | 1123 | | |



Figure 1. Top- female *Culex*; Bottom – male *Culex*




Figure 2. Number of mosquitoes based on sites

4. Discussion

India is inhabitant to over 393 species and 49 genera of mosquitoes. The present study was conducted between the month of July to December 2022 where *Aedes*, *Anopheles* and *Culex* were considered to be most important because of their potential ability in transmission of disease [13].

The present study supports the work of Sule [16] that shows the presence of 393 species of *Culex* which is much higher than that of other genera in Maharashtra. The present work also supports the reported 877 cases of filariasis from the month of January to June 2020 published in The Times of India [13]. Ataullah, studied the diversity of species and relative abundance of *Anopheles*, *Aedes* and *Culex* species in Pakistan and found that *Culex* is dominating followed by *Aedes* over the others which supports the current research [15] Haq and Singh studied the species diversity of mosquitoes in Uttarakhand and found *Culex*, *Aedes* and *Anopheles* to be dominating over other species [14] supported by the work of Kumar who studied the same in Tamil Nadu and found similar results [17]. While studying the abundance of mosquitoes in Jalna [6], it was found that *Anopheles* genus to dominate whereas in the present study where *Culex* was found to be dominating followed by *Anopheles* genus. It was studied that the diversity of mosquitoes from Coimbatore region of Tamil Nadu and found 4 genera of mosquitoes to be predominant, out of which *Culex* was the leading one [18] which further supports our current study. Our work is in sync with the research conducted at Nanded district of Maharashtra wherein *Culex* was found in abundance [2]. The breeding site of *Anopheles* mosquitoes is stagnant water in pond or lakes that are more than half inch in depth whereas the breeding sites of *Culex* and *Aedes* mosquitoes include small puddles or any places for water less than half inch deep [16].

The samples were collected from the places that were habitable by humans such as Rai (Site 1), Navgarh road (Site 2), MBMC Dumping ground (Site 3) and Uttan (Site 4). These sites shows an appropriate breeding place for *Culex* and *Aedes* mosquitoes. It was observed that the number of mosquitoes were higher in the month of September in Nanded District of Maharashtra [4]. A similar study was conducted in United Kingdoms which stated the similar results about the population of mosquitoes being highest in the month of September when the temperature of UK was the highest [17]. But for the present study it was found that the highest number of mosquitoes were found in the month of November and the lowest in the month of July 2022. Mosquitoes are poikilotherms and their temperature is essentially the same as their surroundings, thus they are not capable of regulating their body heat. Mosquitoes are most active at 80°F, become lethargic at 60°F, and cannot function below 50°F [18]. In tropical areas, mosquitoes are active throughout the year. In temperate climates, adult mosquitoes of some species become inactive with the beginning of cool weather and enter hibernation to live through the winter states Central Mass Mosquito Control Project (CMMCP). The temperature in the district of Thane is been fluctuating throughout the year, thus the temperature range in the district was above 29°C in the month of November which is an ideal temperature for mosquitors to remain active and reproduce [5]

For the first time, extensive studies on diversity of mosquitoes in Mira Bhayander have been undertaken and it provides a raw data on the diversity of mosquitoes in this area.

Conclusion

From the present study it can be concluded that there is an increase in the population of *Culex* followed by *Aedes* mosquitoes which is closer to human habitats. With the increasing cases of mosquito borne diseases, there is a need to control the growth of mosquitoes before Filariosis and Dengue become endemic to the area and the city further on. There is also a need to reduce the breeding sites of mosquitoes growth and to increase the awareness regarding the disease and educate on the basic hygiene among the people. The government needs to take standard measures for the same and people need to be thoroughly educated on the matter. The diversity of mosquitoes not only depends on the climate but also on the temperature that they thrive in. They require warm temperatures of approximately 29- 30 °C to flourish well.

Conference disclaimer:

Some part of this manuscript was previously presented and published in the conference: An International Conference on Coastal and Marine Conservation CMC-2024 dated from 1st and 2nd March, 2024 in Mumbai, India. Web Link of the proceeding: <https://mithibai.ac.in/wp-content/uploads/2024/02/CMC2024-CONFERENCE-brochure..pdf>

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.**
- 2.**
- 3.**

References

- [1] Beerntsen, B. T., James, A. A., & Christensen, B. M. (2000). *Microbiology and molecular biology reviews*, 64(1), 115-137.
- [2] Ewing, D. A., Cobbold, C. A., Purse, B. V., Nunn, M. A., & White, S. M. (2016). Modelling the effect of temperature on the seasonal population dynamics of temperate mosquitoes. *Journal of theoretical biology*, 400, 65-79.
- [3] Prakash, A., Jambulingam, P., Rajavel, A., Bhattacharyya, D., Mahanta, J., & Mohapatra, P. (2014). Faunal richness and the checklist of Indian mosquitoes (Diptera: Culicidae). *Check List*, 10(6), 1342-1358.
- [4] Vector borne diseases (2022). World Health Organization. Retrived from <https://www.who.int/news-room/fact-sheets/detail/vector-borne-diseases>
- [5] Dudhmal. D., Chavan. S. and Chinchole. D. (2015). Mosquito Diversity in a Filaria Endemic Village of Nanded District. *Ecology and Fisheries*. 8(1):83-88.

- [6] Jaid, E. L., Nikam, S. V., More, B. V., & Bhandari, J. C. (2011). Diversity of mosquitoes in Jalna urban, Maharashtra state, India. *Journal of Experimental Sciences*, 2(4), 11-12.
- [7] Babu, S., & Nutman, T. B. (2014). Immunology of lymphatic filariasis. *Parasite immunology*, 36(8), 338-346.
- [8] Vogt, M. B., Lahon, A., Arya, R. P., Kneubehl, A. R., Clinton, J. L. S., Paust, S., & Rico-Hesse, R. (2018). Mosquito saliva alone has profound effects on the human immune system. *PLoS neglected tropical diseases*, 12(5), e0006439.
- [9] Tyagi, B. K., Munirathinam, A., & Venkatesh, A. (2015). A catalogue of Indian mosquitoes. *International Journal of Mosquito Research*, 2(2), 50-97.
- [10] Schneider, B. S., & Higgs, S. (2008). The enhancement of arbovirus transmission and disease by mosquito saliva is associated with modulation of the host immune response. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 102(5), 400-408.
- [11] Khan, S. A., Webb, C. E., & Kassim, N. F. A. (2021). Prioritizing mosquito-borne diseases during and after the COVID-19 pandemic. *Western Pacific Surveillance and Response Journal: WPSAR*, 12(2), 40.
- [12] Mira-Bhayander Municipal Corporation City Population Census (2011). Retrieved from <https://www.census2011.co.in/data/town/802786-mira-bhayander-maharashtra.html>
- [13] Maharashtra's Filaria Burden Rises, 877 Cases Recorded Till June (2018). The Times of India. Retrieved from <https://timesofindia.indiatimes.com/city/pune/maharashtras-filaria-burden-rises-877-cases-recorded-till-june/articleshow/6520729.cms>
- [14] Haq I.U. and Singh. S. (2021). Diversity of Mosquito Fauna in different Habitats of District Dehradun, Uttarakhand. *International Journal of Current Microbiology and Applied Sciences*, 10(04): 583-589.
- [15] Attaullah, M., Gul, S., Bibi, D., Andaleeb, A., Ilahi, I., Siraj, M., ... & Ullah, Z. (2021). Diversity, distribution and relative abundance of the mosquito fauna (Diptera: Culicidae) of Malakand and Dir Lower, Pakistan. *Brazilian Journal of Biology*, 83.
- [16] Sule. S. R., Chavan. R. J., Shinde. L. V. and Khaire. B. S. 2021. Climate change, mosquito diversity and epidemics of mosquito borne diseases in India. *Journal of Global Biosciences* Vol. 10(3), pp. 8389- 8403.

- [17] Kumar, K. R., & Nattuthurai, N. (2011). Diversity of mosquito fauna in three selected sites of athoor taluk, Dindigul district, TamilNadu. *Elixir Bio Diversity*, 38, 4057-4059.
- [18] Manimegalai, K. (2010). Studies on the mosquito populations from Coimbatore, Tamil Nadu, India. *Journal of Threatened Taxa*, 961-969.
- [19] Hamza, A. M., & Rayah, E. A. E. (2016). A qualitative evidence of the breeding sites of *Anopheles arabiensis* Patton (Diptera: Culicidae) in and around Kassala town, eastern Sudan. *International journal of insect science*, 8, IJIS-S40071.
- [20] Evans. C. 2019. Mosquito Identification and Key Characteristics of Local Species. Bureau of Environmental Health Services Columbia, South Carolina.
- [21] Agyekum, T. P., Botwe, P. K., Arko-Mensah, J., Issah, I., Acquah, A. A., Hogarh, J. N., ... & Fobil, J. N. (2021). A systematic review of the effects of temperature on *Anopheles* mosquito development and survival: implications for malaria control in a future warmer climate. *International Journal of Environmental Research and Public Health*, 18(14), 7255.
- [22] Bashar, K., Rahman, M. S., Nodi, I. J., & Howlader, A. J. (2016). Species composition and habitat characterization of mosquito (Diptera: Culicidae) larvae in semi-urban areas of Dhaka, Bangladesh. *Pathogens and Global health*, 110(2), 48-61.
- [23] Luo, Q. (1993). A study on transmission of dengue virus by *Culex fatigans*. *Zhonghua liu Xing Bing xue za zhi= Zhonghua Liuxingbingxue Zazhi*, 14(2), 67-69.
- [24] Couper, L. I., Farner, J. E., Caldwell, J. M., Childs, M. L., Harris, M. J., Kirk, D. G., ... & Mordecai, E. A. (2021). How will mosquitoes adapt to climate warming?. *Elife*, 10, e69630.
- [25] McBride, C. S., Baier, F., Omondi, A. B., Spitzer, S. A., Lutomia, J., Sang, R., ... & Vosshall, L. B. (2014). Evolution of mosquito preference for humans linked to an odorant receptor. *Nature*, 515(7526), 222-227.
- [26] Nagpal. B. N. and Sharma. V. N. 1995. Indian *Anophelines*. Oxford & IBH Publication CO. PVT. LTD.
- [27] Rueda, L. M. (2004). *Pictorial keys for the identification of mosquitoes (Diptera: Culicidae) associated with dengue virus transmission*. Walter Reed Army Inst Of Research Washington Dc Department Of Entomology.
- [28] Lee, H., Halverson, S., & Ezinwa, N. (2018). Mosquito-borne diseases. *Primary Care: Clinics in Office Practice*, 45(3), 393-407.
- [29] Andreadis, T. G. (2005). *Identification guide to the mosquitoes of Connecticut* (No. 966). Connecticut Agricultural Experiment Station.

[30] Scheel. M. D., Eggleston. K. K., Achee. N. L., Grieco. J. P., Hapairai. L. 2018. Mosquito control practices and perceptions: An analysis of economic stakeholders during the Zika epidemic in Belize, Central America. PLOS ONE. 13(7).

[31] Amini, M., Hanafi-Bojd, A. A., Aghapour, A. A., & Chavshin, A. R. (2020). Larval habitats and species diversity of mosquitoes (Diptera: Culicidae) in West Azerbaijan Province, Northwestern Iran. *BMC ecology*, 20, 1-11.

[32] Littig. K. S. and Stojanovich. C. J. 2019. Mosquitoes: Characteristics of *Anophelines* and Culicines.

33. Möhlmann TW, Wennergren U, Tälle M, Favia G, Damiani C, Bracchetti L, Koenraadt CJ. Community analysis of the abundance and diversity of mosquito species (Diptera: Culicidae) in three European countries at different latitudes. *Parasites & vectors*. 2017 Dec;10:1-2.

34. Norris DE. Mosquito-borne diseases as a consequence of land use change. *EcoHealth*. 2004 Mar;1(1):19-24.

UNDER PEER REVIEW