

# MORPHOMETRIC ANALYSIS AND CONDITION FACTOR OF FRESHWATER PRAWN, *Macrobrachium lamarrei* (CRUSTACEA - DECAPODA) FROM THE GOMTI RIVER, LUCKNOW (U.P.) INDIA

## ABSTRACT

The present investigations on morphometric analysis and condition factor of freshwater prawn, *Macrobrachium lamarrei* (in both sexes) from The Gomti river, Lucknow, Uttar Pradesh, India. The freshwater prawn is locally known as “Jhinga” was the subject of the study. Prawn samples were collected in different months of both sexes and a close relation between their length and weight were observed. The morphometric analysis showed that females ranged from 31-57mm in length, with their weight ranges from 0.31-1.39g, larger than the males (21-36mm) with their higher weight range (0.13- 0.49g). A high correlation coefficient was observed in males ( $R^2 = 0.808$ ) and females ( $R^2 = 0.881$ ) separately. There were significant differences in the condition factors of males and females at  $P = .05$ . The lowest condition factor (CF) value was observed in January (CF= 1.06) and highest in September (CF= 1.52) in males while in females CF value highest in January (CF=0.82) and lowest in March (CF=0.75). The sex ratio of male: female (1.0 : 2.10) was found, indicating that females significantly greater in number than males.

**Keywords:** Freshwater prawn, *Macrobrachium sp.*, Length-weight relationship, Morphometric analysis, condition factor, The Gomti river

## 1. INTRODUCTION

Crustaceans are important component of aquatic ecosystem. The shrimps, prawns, lobsters and crabs are the good source of protein for humans [14]. Crustacean Fisheries is an important growing industry of the world. Now a days more and more emphasis has been given to freshwater prawn culture across the globe [41, 51]. Similar freshwater prawns like *Macrobrachium lamarrei* and *Macrobrachium dayanum* an “untapped resource”, are promising candidate for aquaculture having various advantages over marine prawn species as well as “Scampy” *Macrobrachium rosenbergii* [56].

The examination of the length- weight relationship plays a crucial role in the field of prawn biology. This investigation holds practical significance as it enables the evaluation of prawn growth in various environmental conditions [8]. The work on various crustaceans including freshwater prawn has also highlighted the importance of length-weight relationship analysis and condition factor to establish correlations between populations from different areas [43, 46]. Furthermore, elucidating the morphometric analysis in aquatic animals, specifically fish, holds significant importance in fisheries and aquaculture [29]. It aids to determining the mean weight corresponding to a given length and also provides variations in growth rate of a specific species in different environmental conditions [2].

The CF of fish reveals the interactions between biotic and abiotic factors of the fish which reflects overall growth and development of animals [49]. [1] reported that the condition factor also reflects the influence of seasonal variations and habitat changes on the species' robustness. Aqua-culturists have documented the morphometric analysis and condition factor of prawns and fishes as crucial indicators of their growth, development and health [7]. Variability in these key parameters (morphometric analysis and condition factors) for prawns provides insights into the suitability of aquatic ecosystem for prawn strength, standing stock, and their production [52, 3].

**Commented [EM1]:** Restructure this statement into something that explain why this study was conceptualized.

**Commented [EM2]:** Provide short methods on the data gathering.

**Commented [EM3]:** There is no gap being stipulated in the introduction section. What particular problem that this study want to address?

**Commented [EM4]:** Provide citation for this claim.

**Commented [EM5]:** fisheries

**Commented [EM6]:** Provide citation

**Commented [EM7]:** Nowadays


**Commented [EM8]:** What other variable being correlated of CF aside from population?

Prawns and shrimps belonging to the genera *Macrobrachium* and *Panaeus* hold significant value worldwide and are often used to enhance the flavor of food preparations due to their high protein content [15, 39]. India, for instance, boasts a rich diversity of these species, with more than 50 of the 200 known *Macrobrachium* species being reported in the country [25]. The freshwater prawn is a relatively small sized prawn species readily available whole year in local habitats. It matures entirely in freshwater and holds promise for freshwater aquaculture [56, 57, 4, 55, 38]. Additionally, this prawn species serves as a bio-indicator for monitoring of environmental pollution [53, 54, 37, 36, 35, 34, 56, 60, 4].

Therefore, the present investigation focused on both morphometric and condition factor analysis of *M. lamarrei* from the river Gomti, Lucknow (U.P.), India.

## 2. MATERIALS AND METHODS

Freshwater prawns, *Macrobrachium lamarrei* were collected from different locations of the river Gomti, Lucknow (U.P.) with the help of local fisherman and brought to the laboratory (N-26°49' 54" E- 80°55'58") and maintained in glass aquaria [53].



map 1 : Collection Site the Gomti river Lucknow(U.P)India (N-26°52'27" E- 80°54'42")

**Commented [EM9]:**

The sampling methodology is not specified, including the absence of designated months or specific seasons for data collection by the researchers.

**Commented [EM10]:** Provide the total number of sampling sites in your study. Provide also description of each sampling site, including the specific coordinates.

**Commented [EM11]:** Provide source and software used in making the map.



Fig 1 : Freshwater prawn, *M. lamarrei*

The total length of each individual was measured with the help of divider, meter scale and vernier caliper and weight by digital weighing machine [ 33, 5, 24, 28, 23] . The male and female prawns were analyzed separately by collecting them according to their gender and by the month for body mass and length-weight relationship [10] .

The length-weight relationship was calculated by linear regression [47] according to the following formula [32].

$$W = aL^b$$

Where :

W= Weight of Prawn(g)

L= Length of Prawn (cm)

a = Intercept ; b= Regression coefficient

The parameters a and b were evaluated by linear regression [ 32, 47, 58, 27, 30] . This equation is sometimes also referred as the length-weight key [12] .

#### Condition Factor(CF)

It shows interaction between exogenous and endogenous factors which reflects the overall growth and development of small prawns. It was calculated using the following formula [17] :

$$CF = 100W/L^3$$

Where :

CF= Condition Factor

Commented [EM12]: Edit this one using the equation format

Commented [EM13]: Edit this one using the equation format

W is the weight (g) of the prawn and L is the Total length of the prawn (cm)

### Sex ratio

By inspecting the presence or absence of appendix masculine in second abdominal pleopod, the sex of the specimens was ascertained. The ratio of male and female was based on the frequency of the prawns caught during the sampling period. The sex ratio of the male and female individuals was calculated [48] .

Sex ratio= M/F,

Where :

M = Number of male prawns

F = Number of female prawns

### Statistical Analysis

The Chi square test was employed to assess the significant difference of sexes every month. All statistical analysis were done using Microsoft Excel 2019 and MINITAB Software.

**Commented [EM14]:** Since the study provide correlation and t-test results but there is no formula and specific test used for correlation and t-test.

## 3. RESULT AND DISCUSSION

The Prawn samples were collected in different months of the both sexes to explain the relationship between their morphometric data . A total of 100 prawns were taken as a sample size. The females were found throughout the year, but the males were not found in the February and March stocks. The males were found in lower numbers compared to female.

**Commented [EM15]:** This is part in the methods sections

A strong positive significant relationship was observed between length and weight of *Macrobrachium lamarrei* ( $R^2 = 0.881$ ) in July, and insignificant relationship was observed in the month of October ( $R^2 = 0.356$ ). But in the males, the relationship was significant in October ( $R^2 = 0.808$ ) and weak relationship observed in the month of September ( $R^2 = 0.147$ ). Table 1 showed morphometric analysis between body mass weight and body length of female freshwater prawn, *Macrobrachium lamarrei* in different months and Table 2 showed the morphometric relationship of the male prawns. The Sex ratio (male and female) of the prawns showed significant difference in various months at ( $P = .05$ ). Therefore the result showed that the sexual frequency and sex ratio varied in different months. The highest sex ratio was observed in the month of September and lowest in February and March (Table 3).

**Commented [EM16]:** Provide implications for this finding. Provide literature findings to support your claim.

**Commented [EM17]:** Provide specific figure of the number males and females.

**Commented [EM18]:** Provide implications of why this is happening

**Commented [EM19]:** Specify the months

The condition factor of the males showed maximum in September (CF = 1.52) and minimum in the month of January (CF= 1.06) ( Table 2) whereas, in the females, condition factor was highest in the January (CF= 0.83 ) and lowest in the March ( CF= 0.75) ( Table 1) . A positive allometric growth of *Macrobrachium lamarrei* from the Gomti river on the basis of growth coefficient (b value) by using t-test (P= .05) (Table 1 and 2) . The values of growth coefficient were high in the month of September , which indicates the suitability of the environment for food availability in the Gomti river.

The study of the length-weight relationship of *M. lamarrei* showed a highly significant correlation between total length and body weight. The present study showed, the regression coefficient (b) for females is moderately high as compared to the males. This suggests that the females gain more weight with increase in the length, indicating better well-being. [9] , also reported same observations in freshwater minor carp fish, *Puntius conchoni* from the Mandal river. The regression coefficient values showed a positive allometric growth pattern, which corresponds to the outcomes of [45, 6] regarding the *M. macrobrachion*, and [31] reported in *P. notialis*. Moselen & Miebaka., 2018, suggests that prawns become relatively robust and stronger as they increased in length when showing positive allometric growth patterns. Strong regression coefficient between length and weight were observed, with coefficients for males (R2 = 0.14-0.80) and for females (R2 = 0.35-0.88), indicating that the model is suitable for the study.

Table 1: Regression analysis and condition factor of the Female freshwater prawn, *Macrobrachium lamarrei*

Month	Weight (g) ± SE	Length (mm) ± SE	R <sup>2</sup>	b	CF Value
-------	-----------------	------------------	----------------	---	----------

January	0.83 ± 0.09	45.7 ± 0.13	0.768	1.25	0.83
February	0.81 ± 0.05	46.9 ± 0.10	0.528	1.37	0.78
March	0.58 ± 0.03	42.5 ± 0.08	0.635	2.09	0.75
April	0.83 ± 0.08	46.2 ± 0.16	0.815	1.76	0.81
May	0.69 ± 0.06	44.7 ± 0.16	0.761	2.14	0.76
June	0.89 ± 0.07	47.7 ± 0.11	0.366	0.87	0.81
July	0.76 ± 0.05	45.2 ± 0.12	0.881	2.06	0.81
August	0.83 ± 0.08	46.7 ± 0.14	0.660	1.37	0.79
September	0.88 ± 0.07	46.9 ± 0.11	0.812	1.39	0.82
October	0.73 ± 0.05	45.8 ± 0.12	0.356	1.34	0.77
November	0.87 ± 0.06	47.1 ± 0.11	0.806	1.53	0.81
December	0.77 ± 0.06	45.5 ± 0.11	0.803	1.63	0.80

Table 2: Regression analysis and condition factor of the Male freshwater prawn, *Macrobrachium lamarrei*

Month	Weight (g) ± SE	Length (mm) ± SE	R <sup>2</sup>	b	CF Value
January	0.25 ± 0.01	28.7 ± 0.07	0.730	4.49	1.06
February	-	-	-	-	-
March	-	-	-	-	-
April	0.27 ± 0.02	27.8 ± 0.06	0.500	1.93	1.27
May	0.31 ± 0.02	28.4 ± 0.06	0.549	2.11	1.37
June	0.27 ± 0.07	27.9 ± 0.04	0.765	5.10	1.25
July	0.27 ± 0.07	27.8 ± 0.05	0.695	5.61	1.29
August	0.23 ± 0.01	26.7 ± 0.08	0.709	4.55	1.23
September	0.30 ± 0.01	27.5 ± 0.07	0.147	1.41	1.52
October	0.25 ± 0.00	25.9 ± 0.05	0.808	6.52	1.45
November	0.29 ± 0.02	28.2 ± 0.06	0.578	1.82	1.27
December	0.24 ± 0.01	25.9 ± 0.08	0.819	5.88	1.40

Commented [EM20]: Why there is no data in these months?

Table 3: Sex ratio of Male and Female freshwater prawn, *Macrobrachium lamarrei*

--	--	--

<b>Month</b>	<b>M/F Ratio</b>	<b>GDC</b>
<b>January</b>	0.28	2
<b>February</b>	0	100
<b>March</b>	0	100
<b>April</b>	0.36	1
<b>May</b>	0.25	20
<b>June</b>	0.31	4
<b>July</b>	0.28	2
<b>August</b>	0.42	10
<b>September</b>	0.47	4
<b>October</b>	0.38	4
<b>November</b>	0.28	2
<b>December</b>	0.31	4

According to the factors such as food availability, feeding rate and reproductive activity the length-weight relationship may temporarily vary, however, the parameter 'b' usually insignificant throughout the year [11]. In *M. armatus*, the regression coefficient showed gender diversity for male and female fishes as reported by [50].

[26] reported length-weight relationship of *Macrobrachium vollehoveni* with regression coefficient value ranged from 2.58 to 3.11 from Badagry creek. The analysis of length-weight relationship was given by [32] in *Perca fluviatilis*, showed the cubic parabola into general parabola. [22, 40], inspected the value of 'b' varying from 2.5 to 4.0. The evaluated parameter 'b' revealed negative allometric growth for all the species and was supported by high correlation values, suggesting elevated predictability between carapace length and weight of the studied shrimps [13, 18].

In the present study, *Macrobrachium lamarrei* reflected the highest condition factor for the males (1.53 in September) and the females (0.83 in January). There were notable differences between male and female in different months (April, June, July, September) of the year. Similar results were observed by [5] in *M. nipponense* in the Siah Darvishan River, [16] in *Metapenaeus monoceros*, Bhidiya and [6], in *M. macrobrachion* from the Kwa-river.

[28] reported Fulton's condition factor for *Labeo rohita* which ranged from 0.77506-2.45703. The key factors which affect the K value consists of size, season, sex and development of gonads in fish [21]. The well-being of the fish is also influenced by food supply, ecological factors etc. [32]. [59, 20, 19] reported that the value of K tends to decline with age. In Southern coasts of the Caspian Sea, Iran, condition factor of *M. nipponense* were found 0.89-1.90 [44].

The gender distribution in the present studies varies significantly from month to month. February and March show a notable gender imbalance with 0 males and 100 females each. This suggests that these months might represent a specific stage in the prawn's life cycle or behavior, such as reproduction or molting when males are less active or not present.

The presence of 100 females in February and March indicates that these are the months when females release their eggs or juveniles. The Greatest common divisor (GCD) provides information about some characteristic of the prawn population, possibly related to reproduction. For example, a GCD of 100 prawns in February and March might indicate that these are peak breeding months when the entire population consists of females (e.g., berried females). The sex ratio varies across the months and is affected by the GCD.

In months with a GCD of 100 prawns, the ratio is 0:1, indicating a complete absence of males compared to females. This reinforces the idea that these months are associated with reproductive activities of the female prawns.

In contrast, months with lower GCDs (e.g., in June, September, October, and December) have male-to-female ratios ranging from 6:19 to 7:18, indicating a more balanced or slightly female-biased population.

#### 4. CONCLUSION

*Macrobrachium lamarrei* shows a positive allometric growth in the Gomti river. The elevated value of regression coefficient "R" showed a strong relationship between total length and body weight. The high value of the coefficient of determination R<sup>2</sup> suggested that the model

used for the analysis fits the data, hence confirms the fitness of the model. Knowledge of morphometric analysis and the condition factor of introduced or invaded species are essential to assess an appropriate management of alien and native species in an aquatic ecosystem. More advance studies are required to determine where there is a correlation between each morphotype's physiological, behavioural and functional characteristics. An understandable explanation of morphotype of this species is very important for understanding growth processes and adaptive value in the population to improve culture management.

Commented [EM21]: What model?

## REFERENCES

1. Abowei, J.F.N. The condition factor, length -weight relationship and abundance of *Ilisha africana* (Block, 1795) from Nkoro River Niger Delta, Nigeria. *Adv. J. Food Sci. Technol*, 2010. 2(1): 6-11.
2. Abowei, J.F.N.; Tawari, C.; Cdeekae, S.N. and Amakiri, N.E. A study of the length – weight relationship and condition factor of *Pseudotolithus elangatus* 559 Fulton's Condition Factor of *Macrobrachium nipponense* in Siah Darvishan River (Browdich, 1825) from Bonny Estuary, Niger Delta, Nigeria. *Int. J. Trop. Agr. Food.Sys*, 2008. 2(3-4): 249-254.
3. Adeosun FI, Adedoyin IA, Oghenebrorhie OMT, Quddus AA, Micheal AO. Morphometric and Meristic features and Length-weight Relationship as indicators of quality of *Brycinus macrolepidotus* in Lower River Ogun, Nigeria. *Egyptian Journal of Aquatic Biology & Fisheries.*; 2019. 23(2):433-441.
4. Ahmed. A., Lodhi. S. and Shukla. S. Observations on feeding behaviour of fresh water prawn, *Macrobrachium lamarrei* (Crustacea: Decapoda) *Int. J. of Fish. and Aqua. Stud*. 2021. 9(6):109-112.
5. Aminisarteshnizi. M. Length -weight relationship and Fulton's Condition Factor of *Macrobrachium nipponense* (De Haan, 1894) in Siah Darvishan River, Iran. *Egyptian Journal of Aquatic Biology and Fisheries.*,2021. 25(2): 551-560.
6. Andem, A. B.; Idung, J. U.; George, E. and Ubong G.U. Length-weight relationship and Fulton's Condition Factor of brackish river prawn (*Macrobrachium macrobrachion*, Herklots, 1851) from Great Kwa River, Obufa Esuk beach, Cross River state, Nigeria. *European Journal of Experimental Biology*, 2013. 3(3):722-730.
7. Attia AO, El-Aiatt, Kariman A. Sh. Shalloof. Study on the biology of the small scaled terapon *Terapon puta* (Cuvier, 1829) from Bardawil lagoon, North Sinai, Egypt. *Egyptian Journal of Aquatic Biology & Fisheries*. 2019. 23(2):95 – 107.
8. Bahuguna P, Joshi HK. Statistical observation on the length-weight relationship of brain and body in a coldwater catfish *Amblyceps mangois* (Ham. -Buch.) from Garhwal region. *Environ. Conser. Journal*. 2010;11(1-2): 21-23.
9. Bahuguna P, Shah KK, Kumar, R. Observation on the length-weight relationship and relative condition factor of *Barilius Bendelisis* (Ham.) inhabiting a spring-fed tributary of river Alaknanda (Garhwal Himalaya), India. *J.Natcon*. 2009;21(2):215-220.
10. Bahuguna. P., Dimri. A., Rayal. R. and Sharma. N. Observations on the body mass weight-length relationship and relative condition factor of *Macrobrachium assamensis peninsularis* (Tiwari) from Khoh river, Uttarakhand, India. *Uttar Pradesh journal of Zoology*. 2021. 43(13): 54-65.
11. Begenal T.B, Tesch F.W. Age and growth in methods of assessment of fish production in freshwater, Ed. Bagenal, T. Oxford. Blackwell Scientific Publication., 1978. 101-136.

12. Biswas S.P. Length-weight relationship and condition factor. In: Manual of methods in Fish Biology. South Asian Publishers, New Delhi. 1993. p. 60-64.
13. Carlander K.D. Handbook of freshwater fishery biology. Ames. Fishing News Books., 1969. 1.
14. Carnevali R.P., Collins P.A. and Poi de Neiff A.S.G. Trophic ecology of the freshwater prawn, *Pseudopalaemon bouvieri* (Decapoda: Palamonidae) in Northeastern Argentina, with remarks on population structure. Rev. Biol. Trop. 2012. 60 (1): 305-316.
15. Cruz, Y.M. and Olatunbosun, O. Comparative Study on the efficiency of three different types of crab pot in Iceland fishing ground. UNU-Fisheries Training Program, 2013. 4:1-25.
16. Dineshbabu, A.P. Length-weight relationship and growth of the speckled shrimp *Metapenaeus monoceros* (Fabricius) off Saurashtra. Journal of Marine Biology Ass. India, 2006. 48 (2): 180 - 184.
17. Enin U. Length- weight parameters and condition factor of two West African Prawns. Rev. Hydrobiol. Trop., 1994. 27: 121-127.
18. Froese R. Cube Law. Condition factor and weight-length relationship: History, meta-analysis and recommendations. J. Appl. Ichthyol., 2006. 22:241-253.
19. Gandotra R., Vivek., Singh. D and Shanker R. Length-weight relationship and condition factor of *Aspidoparia morar* (Ham.) inhabiting river Tawi and its tributaries. Biosciences, Biotechnology Research Asia., 2009. 6(2): 767-772.
20. Gandotra. R., Shanker R., Ahmed S. and Sagar S. Morphometry of *Tor putitora* (Ham.) from Jhajar stream, Jammu (J&K). J. Inland Fish Soc. India., 2008. 40(1): 86-89.
21. Heincke F. Bericht über die Untersuchungen der Biologischen Anstalt auf Helgoland zur Naturgeschichte der Nutzfische. (1. April 1905 bis 1. Oktober 1907). [Report on the investigations of the Biological Institute on Helgoland on the natural history of commercial fish (April 1, 1905 to October 1, 1907). In: Die Beteiligung Deutschlands an der Internationalen Meeresforschung, 4. & 5. Jahresbericht. Verlag von Otto Salle, Berlin; c1908.
22. Hile R. Age and growth of the Cisco, *leucichthys artedti* (Le seum) in the North-Eastern high lands. Wisconsin Bull U.S. bur. Fish., 1936. 48:211-317.
23. Ibrahim. S., Zhong. Z., Lan. X., Luo. J., Tang. Q., Xia. Z., Yi. S. and Yang. G. Morphological Diversity of Different Male Morphotypes of Giant Freshwater Prawn *Macrobrachium rosenbergii* (De Man, 1879). Aquaculture Journal., 2023. 3: 133-148.
24. Iloba, Isioma. K, Otokuefor, Zino. J and Israel. O.S. Length-weight relationship and health of *Macrobrachium dux* (Lenz, 1910) and *Macrobrachium macrobrachion* (Herklots, 1851) in river Ethiop Headwaters, Nigeria. Uttar Pradesh Journal of Zoology., 2022. 43(5): 1-13.
25. Jayachandran K.V., and Indira. Prawn Fisheries Resources of India for food security as well as for rural employment. 97th Indian Sci. Congr..Symposium, 2010. January 3-7, Thiruvananthapuram.
26. Jimoh A.A., Fakoya K.A., Hamed A.M., Amosu A.O. and Kumolu- Johnson C.A. Meristic and Morphometrics in the African River Prawn, *Macrobrachium vollenhovenii* (Herklots, 1857) from Ologe Lagoon, South West, Nigeria. Journal of Agriculture and Environmental Research Studies. 2005, Vol 1(1); 12-18.
27. Jones R.E., Petrell R.J. and Pauly D. Using modified length-weight relationships to assess the condition of fish. Aquacultural Engineering, 1999. 20: 261-276.

28. Kanaujia. S., Sahu. A., Verma D.K. and Singh. M.B. Length-weight relationship and condition factor of a commercial food fish *Labeo rohita* (Hamilton, 1822) from Ghaghara River, Uttar Pradesh, International Journal of Fisheries and Aquatic Studies., 2023. 11(4): 96-99.
29. Khanipour, A.A.; Noori, A.; Amini, M. and Kamrani, E. Length-weight relationship and Fulton's condition factor of *Macrobrachium nipponense* (De Haan, 1849) in Anzali lagoon of Iran. Iranian Journal of Fisheries Sciences, 2020. 19(1): 496-500.
30. Koutrakis E.T. and Tsikliras A. C. Length- weight relationships of fishes from three northern Aegean estuarine systems (Greece). J. Appl. Ichthyol. 2003. 19: 258-260.
31. Lawal-Are A.O. and Akinjogunla V. F. Length- Weight Relationships and Condition Factor in Lagos Lagoon, South West, Nigeria. Science and Technology, 2012. 2(3): 32-40.
32. Le Cren ED. The length-weight relationship and seasonal cycle in gonad weight and condition in Perch (*Perca fluviatilis*). J. Anim. Ecol. 1951. 20:201-219.
33. Lima J.D.F., Da Cruz M.C.M. and Silva L.M.A.D. Reproductive biology of *Macrobrachium surinamicum* (Decapoda: Palaemonidae) in the Amazon River mouth. Acta Amazonica., 2015. 45(3): 299 – 306.
34. Lodhi H. S., Shukla. S. and Sharma U.D. Studies on structure of heart of two freshwater prawns under light microscope. Uttar Pradesh J. Zool. 2009. 29(1):1-10.
35. Lodhi H. S., Tiwari K. J., Shukla.S. and Sharma U.D. Copper induced fluctuations in total haemocyte counts (THCs) of freshwater prawn, *Macrobrachium lamarrei* (Crustacea- Decapoda). J. Env. Bio. Sci. 2008. 22(2):135-142.
36. Lodhi H.S., Khan M.A., Verma R. S. and Sharma U. D. Seasonal fluctuations in the total haemocyte counts (THC's) of freshwater prawn, Him. J Env. Zool. 2004. 18(1):1-5.
37. Lodhi. H.S., Khan. M.A., Verma R.S. and Sharma U.D. Acute toxicity of Copper Sulphonate to freshwater prawns. J. Environ. Biol. 2006. 27(3):585-588.
38. Lodhi. S., Ahmed. A., Lodhi.H.S. and Shukla. S. A report on freshwater prawn, *Macrobrachium lamarrei* (Crustacea: Decapoda) from river Gomti, Lucknow (U.P.), India. Journal of Experimental Zoology. India, 2024. Vol 27(1): 751-756.
39. Major, R.N., Taylor, D.I. Conner, S. and Conner G. Factors affecting bycatch in a developing New Zealand scampi potting fishery. Fisheries Research, 2016. 10:55-64.
40. Martin W.R. The mechanics of environmental control of body form in fishes. Univ. Toronto Stud. Biol. 58: Publ. Ant. Fish Res. Lab., 1949. 70:1-91.
41. Mohanty K.R. and Sunderray K.J. Environmental impacts of coastal shrimp farming and measures to reduce consequential pollution. Fishing Chimes, 1999. 19: 7.
42. Moslen, M. and Miebaka, C. A. Condition factor and length-weight relationship of two estuarine shell fish (*Callinectes* Sp and *Penaeus* sp) from the Niger Delta, Nigeria. International Journal of Fisheries and Aquatic Studies, 2018. 6(1): 188-194.
43. Mossolin, E. C. & Bueno, S. L. S. Relative growth of the second pereiopod in *Macrobrachium olfersi* (Wiegmann, 1836) (Decapoda, Palaemonidae). Crustaceana 2003. 76(3):363-376.
44. Namin, J.I.; Nami, E. and Heidary, S. Length-Weight Relationship and Fulton's Condition Factor of *Macrobrachium nipponense* (Dehaan, 1849) in southern coasts of the Caspian Sea-Iran. International journal of Advanced Biological and Biomedical Research, 2014. 2(5):1650-1656.
45. New M.B., Valenti W.C., Tidwell I.H., D'Abram.o.L., and Kuttey M.N. Freshwater prawns: Biology and farming . 2010. Willy – Blackwell Publishing Ltd.

46. Pantaleão, J. A. F.; Hirose, G. L. & Costa, R. C. Relative growth, morphological sexual maturity, and size of *Macrobrachium amazonicum* (Heller 1862) (Crustacea, Decapoda, Palaemonidae) in a population with an entirely freshwater life cycle. *Invertebrate Reproduction & Development*. 2012. 56(3):180-190.
47. Pauly D. Some simple methods for the assessment of tropical fish stocks. *FAO fish tech Rap*, 1983. 234.52.
48. Peña-Mendoza, B.; Gómez-Márquez, J.L.; Salgado-Ugarte, I.H. and Ramírez-Noguera, D. Reproductive biology of *Oreochromis niloticus* (Perciformes: Cichlidae) at Emiliano Zapata dam, Morelos, Mexico. *Rev. Biol. Trop*, 2015.53 (3-4): 515-522.
49. Rajeevan R., Edirisinghe U. and Athauda ARSB. Length- weight relationships and Condition factor of giant freshwater prawn, *Macrobrachium rosenbergii* (De Man, 1879) in five perennial reservoirs in Northern Province, Sri Lanka. *International Journal of Fisheries and Aquatic Studies*. 2018. 6(5): 283-287.
50. Rashid M, Bahuguna, P, Dobriyal AK. Analysis of length- Weight relation and relative condition factor of the hill stream fish *Mastacembelus armatus* (laceped) from river Nayer, Garhwal, Uttarakhand. *Int. J. Recent Sci.Res*. 2019;10(01): 30574-30580. (January).
51. Sankar G, Elavarasi A, Sakkaravarthi K, Ramamoorthy K. Biochemical changes and growth performance of Black tiger Shrimp larvae after using *Ricinus communis* extracts as feed additive. *Intl. J Phar Tech. Res.*,2011. 3; 201-208.
52. Shallof KA. Sh, El-Far AM. Length-Weight Relationship and Condition Factor of Some Fishes from the River Nile in Egypt with Special Reference to Four Tilapia Species. *Egyptian Journal of Aquatic Biology & Fisheries*. 2017. 21(2):33-46.
53. Sharma U.D. and Shukla, S. Behaviour dysfunction of fresh water prawn, *Macrobrachium lamarrei* (Crustacea-Decapoda) following exposure to synthetic detergent, linear alkyl Benzene Sulphonate. *Biol. Mem*. 1990. 16(12):58-61.
54. Sharma U.D. and Shukla. S. Acute toxicity of heavy metals and detergent to fresh water prawn, *Macrobrachium lamarrei* (Crustacea-Decapoda). *Him. J. Env. Zool*. 2006. 20(1):1-6.
55. Sharma. M., Lodhi. S. and Ahmed. A. Freshwater prawn aquaculture: Prospects in U.P. *Int. J. of Fish. and Aqua. Stud*. 2022. 10(4):146-151.
56. Shukla. S. and Sharma U.D. Smaller fresh water prawns: Their aquaculture potential and suitability as good laboratory model. In: *Bioresources for food security and rural livelihood*. Kulkarni G.K. and Pandey P.N. (Eds). 2010. PP.189-204, Narendra Publications, Delhi.
57. Shukla. S., Shukla. R., Shukla. S., Ahmad. A. and Mishra.A. Effect of temperature on chromatophores of freshwater prawn, *Macrobrachium lamarrei* (Crustacea-Decapoda). *Anusandhaan*. 2017. 5 (1):80-84.
58. Sparre. P., Ursin. E. and S.C. Venema. Introduction to tropical fish stock assessment. Part 1. *Manual FAO Fisheries Technical*. FAO Rome, 1989. pp: 337.
59. Tenali. D.R., Phom N.P., Yamala S.P., Gudipudi S.B., Kashyap.S. and Nirmalkar. R. Evaluation of Length-weight relationship and condition factor among the cohorts of *Labeo calbasu* in Pachipenta reservoir, Andhra Pradesh. *Journal of Experimental Zoology.*, 2023. 26(2): 2329-2332.
60. Verma D. R., Lodhi H. S., Tiwari K. J., Shukla. S. and Sharma U. D. Copper sulphate induced changes in scaphognathite oscillations and oxygen consumption of fresh water

prawn, *Macrobrachium lamarrei* (Crustacea-Decapoda). Journal of Applied and Natural Science. 2010. 2(1):34-7.

UNDER PEER REVIEW