

# TAXONOMIC CHECK LIST OF ICHTHYOFAUNAL DIVERSITY IN GOSTHANI RIVER AT TAGARAPUVALASA, VISAKHAPATNAM, ANDHRA PRADESH, INDIA

## ABSTRACT

The current study examined the taxonomy and assortment of fish species in the Gosthani River in Andhra Pradesh from April 2023 to March 2024. A total of 47 fish species were documented at the two landing sites of Gosthani River. The fish samples were carefully cleansed, and photos were taken on location. These fish were fixed in glass jars and kept in a 10% formalin solution. Following that, the samples were identified up to the species level using classical taxonomic techniques. Order Cypriniformes was observed to have the highest contribution to the species diversity: 15.78% families, 38.23% genera, and 40.42% species, followed by Siluriformes (26.31% families, 17.64% genera, and 21.73% species); Belontiiformes (10.52% family, 5.88% genera, and 4.25% species); Cichliformes (1 family, 3 genera and 3 species); Perciformes and Synbranchiiformes (1 family, 2 genera and 2 species); Anguilliformes (1 family, 1 genera and 2 species); Channiformes (5.26% family, 8.82% genera, and 6.38% species); Cyprinodontiformes, Gobiiformes, Osteoglossiformes, and Anabantiformes (5.26% family, 2.94% genera, and 2.12% species). The taxonomic trophic levels were found to have the largest percentage of omnivores have 46.81%, followed by carnivorous 34.04% and herbivorous 19.14%. The composition of taxa, population status, trophic levels, and IUCN status were reported in this research paper.

**Key words:** Ichthyofauna, Population status, Trophic level, Herbivorous, Omnivorous, Carnivorous, IUCN

## INTRODUCTION

The Gosthani River is a medium-sized, east-flowing river in Andhra Pradesh, India. It originates in the Ananthagiri Hills of the Eastern Ghats and flows for 120 kilometers across Thatipudi reservoir before entering the Bay of Bengal at Bheemunipatnam to form an estuary. The Gosthani River is a minor river basin with a total drainage area of less than 2000 km<sup>2</sup>. The depth of the river's deepest point varies by location and 1.5-2 meters in the river's middle part. The Vizianagram and Vishakhapatnam districts rely heavily on the Gosthani River for drinking water. The water from this River has been diverted for commercial and agricultural use. Riverine fisheries are a sort of inland fishing where fish are caught straight from river systems using scientifically designed gear. India has vast riverine fisheries resources due to the presence of several fertile riverine systems.

According to Kar et al., (2003), the Indian subcontinent has around 2500 fish species, 930 of which are freshwater and 1570 of which are marine. There are 801 freshwater fishes present (Fish base 2004). Ichthyofaunal diversity refers to the diversity of fish species that exist depending on context and magnitude; it can relate to alleles or genotypes within life forms within a fish community as well as species or life forms that exist throughout aquaculture environments (Burton et al., 1992). There are approximately 21,723 extant fish species in the world, including 8,411 freshwater species and 11,650 marine forms. India is one of the world's mega-biodiversity countries, ranking ninth in

**Comment [U1]:** This study was designed to examine the

**Comment [U2]:** Increase the sample size at least to give a representative results.

terms of freshwater mega-biodiversity (Mittermeier&Mitemeir, 1997). Biodiversity is the degree of variation of living forms within a particular ecosystem; biodiversity is necessary for ecological stabilization, conservation of overall environmental quality, and comprehending the inherent worth of all species on the planet, as expressed by Ehrlich et al., (1991). Traditional fishers in Gosthani river and surrounding regions, fisher families from Scheduled and Backward Castes, have fishing rights in these bodies of water, which have been recognized by the Panchayats or Fisheries Department as Fishermen Cooperative Societies. The current study provides the most recent database of fish species and is the first documentation of fish fauna in Gosthani River.

**Comment [U3]:** Expand on literature review to bring in more information from previous studies elsewhere.

## MATERIAL AND METHODS

The fish samples were collected out fortnightly from April, 2023 to March 2024 from three landing stations (Thagarapavalasa (17.937486N, 83.420869E). Padmanabham (17.966390N, 83.355637E) Fig 1& 2. Representative species collected carefully with the help of local fishermen by using different types of gears and baskets traps (Rama Rao 2014). In the laboratory, the samples were washed thoroughly and images were captured. These fish were fixed in glass jars before being preserved in a 10% formalin solution. Following that, the samples were recognized up to the species level using classical taxonomic methods such as morphometric features, meristic counts, and descriptive characters. Fish species identification was validated using guidelines produced by Day (1958), Talwar and Jhingran (1991), Jayaram (1999, 2011), Menon 1999, and Munro 2000).




Fig 1. Gosthani River



Fig 2. Godhani River at Tagarapuvalasa and Padmanabham

**Comment [U4]:** Include pictorial steps on how classification was done to further add more clarity to future readers of your publication.

## RESULTS & DISCUSSIONS

The current study assessed the taxa and diversity of fish species in the Godhani River in Andhra Pradesh from April, 2023 to March 2024. A total of 47 fish species were documented in the two selected landing stations. An updated, comprehensive checklist of finfish with their current scientific names, trophic level, fishery status, and IUCN status was prepared after verification with published literature and web-based information such as Froese and Pauly, 2020, (FishBase), Eschmeyer 2023 (Catalog of Fishes) (Table 1). According to CIFE (2011), there are 64 fish species in the Godavari River in Maharashtra, belonging to 15 distinct families and 38 genera, from Gangapur Dam to Raheer. Chinnababu et al., (2021) reported a total of 50 fish species from the selected locations of the Godavari river at Rajamahendravaram, which was significantly fewer than the current documented species.

In the present study, a total of 47 finfish species belonging to 11 orders, 19 families and 34 genera were recorded from Godhani River. Among the 11 orders, the order Cypriniformes was observed to have the highest contribution to the species diversity (15.78% families, 38.23% genera and 40.42% species), followed by Siluriformes (26.31% families, 17.64% genera and 21.73% species); Belontiiformes (10.52% family, 5.88% genera and 4.25% species); Cichliformes (1 family, 3 genera and 3 species), Perciformes and Synbranchiformes (1 family, 2 genera and 2 species); Anguilliformes (1 family, 1 genera and 2 species); Channiformes (5.26% family, 8.82% genera and 6.38% species); Cyprinodontiformes, Gobiiformes, Osteoglossiformes, and Anabantiformes (5.26% family, 2.94% genera and 2.12% species) Table 2; Fig 3. Chandra Sekhara Rao et al., (2014) identified 66 fish species from 9 orders, 22 families, and 38 genera. Order Cypriniformes had the most dominant group with 26 species, followed by Siluriformes and Mugiliformes each with 11 species, Perciformes with 7 species, Anguilliformes, Cyprinodontiformes, and Mastacembeliformes each with 3 species, and Osteoglossiformes and Elopiformes each with 1 species. Rama Rao (2014b) reported at Lower Manair Dam at Karimnagar Dt

The taxonomic trophic levels in the present study was classified into herbivorous (2.0-2.5), omnivore (2.5-3.5), and carnivorous (3.5-4.5) based on their trophic level. The omnivores have a highest percentage of 22 (46.81%), followed by the carnivorous 16 (34.04%), and the herbivorous 09 (19.14%) (Tab. 3, Fig. 4). A similar study was observed by Rama Rao (2023) reported the highest number of omnivores (51.02%, followed by carnivores (26.53% and herbivores (18.36%) in Gotta Barrage at Hiramandalam. The trophic level community structure of recorded fish species

demonstrated the dominance of top-level carnivores (39%), followed by mid-level carnivores (28%), predators (17%), omnivores (14%), and herbivores or planktivores (2%), according to Haojie Su (2021). The majority of the finfish species identified as omnivorous throughout this study. In the present investigation the number and percentage composition of Population Status was 25 species were common which contributed to 53.19%, followed by 11 species were abundant which contributed to 23.40%, 7 species were abundant which contributed to 14.89% and 04 species are moderate which contributed to 08.51% in the total taxa (Table 4, Fig. 5). Rama Rao (2023) the similar results were reported to 25 species were common which contributed to 51.02%, 12 species were abundant which contributed to 24.48%, 7 species are moderate which contributed to 14.28% and 5 species were moderate which contributed to 10.20% in the total catch at Gotta Barrage. Rama Rao (2014b) represented the number and percentage composition of Population Status is 19 species were abundant contributed to 29.69%, 21 species common (32.81%), 14 species moderate (21.86%) and 10 species rare (15.63%) in the total catch at Lower Manair Dam,

According to IUCN status 39 species contributed to 82.97% are least concern (LC), six species contributed to 12.76% are near threaten (NT), one species were not evaluated (NE) and data deficient (DD) with 2.12%, Table 4, Fig 6. The similar study was reported by Priyanka et al., (2021) represents the IUCN red list categories, 52.5% of the species are least concern, 20% are not evaluated, 10% are near threatened, 5% are data deficient, 5% are lower risk near threatened and vulnerable, and 2.5% are lower risk least concern (Rama Rao and Vinod Kumar, 2017). Rama Rao and Ramachandra Rao (2021) observed the Ichthyofaunal diversity at Narayanapuram Anicut at Nagavali River. Further species diversity was found to be the highest during Monsoon season compared to other seasons. Rama Rao (2018) reported more Least concerned fish species at Kalinga Dal reservoir. Rama Rao (2014b) identified 51 species (79.69%) are least concern (LC), 3 species (4.69%) are not evaluated (NT), 4 species (6.25%) are data (EN), vulnerable (VU) and not evaluated (NE) in Lower Manair Dam at Karimnagar Dt. In the present study the ichthyofauna species percentage contribution was nearly similar and *Labeorohita* and *Catlacatla* are found to be dominant species in total catch in the sampling stations.

Table: 1. Taxa of fishes and their order, family, genus, species, population status and IUCN status at Gosthani River

No.	Order / Family	Scientific Name	Trophic level	Population Status	IUCN Status
1	Anguilliformes/ Anguillidae	<i>Anguilla bengalensis</i>	3.8	R	NT
2	Anguilliformes/ Anguillidae	<i>Anguilla bicolor</i>	3.6	R	NT
3	Beloniformes/ Belonidae	<i>Xenentodon cancila</i>	3.9	R	DD
4	Beloniformes/ Hemiramphidae	<i>Hyporhamphus limbatus</i>	3.1	R	LC
5	Cypriniformes/ Cyprinidae	<i>Cirrhinus mrigala</i>	2.4	A	LC
6*	Cypriniformes/ Cyprinidae	<i>Ctenopharyngodon idella</i>	2.0	M	LC
7*	Cypriniformes/ Cyprinidae	<i>Hypophthalmichthys molitrix</i>	2.0	R	NT
8	Cypriniformes/ Cyprinidae	<i>Labeo catla</i>	2.8	A	LC
9	Cypriniformes/ Cyprinidae	<i>Labeo calbasu</i>	2.0	C	LC
10	Cypriniformes/ Cyprinidae	<i>Labeo rohita</i>	2.2	A	LC

11	Cypriniformes/ Cyprinidae	<i>Osteobramacotio</i>	2.9	C	LC
12	Cypriniformes/ Cyprinidae	<i>Puntius chola</i>	2.5	A	LC
13	Cypriniformes/ Cyprinidae	<i>Puntiusticto</i>	2.2	A	LC
14	Cypriniformes/ Cyprinidae	<i>Puntiussophore</i>	2.6	A	LC
15	Cypriniformes/ Cyprinidae	<i>Systemussarana</i>	2.9	C	LC
16	Cypriniformes/ Cyprinidae	<i>Garragotyla</i>	2.0	R	LC
17	Cypriniformes/ Danionidae	<i>Amblypharyngodonmicrolepis</i>	3.3	A	LC
18	Cypriniformes/ Danionidae	<i>Amblypharyngodonmola</i>	3.3	A	LC
19	Cypriniformes/ Danionida	<i>Salmostomabacaila</i>	3.2	C	LC
20	Cypriniformes/ Danionida	<i>Salmostomaphulo</i>	3.2	C	LC
21	Cypriniformes/ Danionida	<i>Rasboradaniconius</i>	3.1	C	LC
22	Cypriniformes/ Danionidae	<i>Daniodevario</i>	3.0	C	LC
23	Cypriniformes / Cobitidae	<i>Lepidocephalichthysguntea</i>	2.7	M	LC
24	Cyprinodontiformes/ Aplocheiidae	<i>Aplocheiluspanchax</i>	3.8	C	LC
25	Channiformes/ Channidae	<i>Channaorientalis</i>	3.8	C	NE
26	Channiformes/ Channidae	<i>Channapanctata</i>	3.8	A	LC
27	Channiformes/ Channidae	<i>Channa striatus</i>	3.6	C	LC
28	Gobiiformes/ Gobiidae	<i>Glossogobiusgiuris</i>	3.7	C	LC
29	Osteoglossiformes/ Notopteridae	<i>Notopterusnotopterus</i>	3.5	C	LC
30	Siluriformes/ Bagridae	<i>Mystusbleekeri</i>	3.3	C	LC
31	Siluriformes/ Bagridae	<i>Mystuscavasius</i>	3.4	C	LC
32	Siluriformes/ Bagridae	<i>Mystusgulio</i>	4.0	R	LC
33	Siluriformes/ Bagridae	<i>Mystustengara</i>	3.2	A	LC
34	Siluriformes/ Bagridae	<i>Mystusvittatus</i>	3.1	A	LC
35	Siluriformes/ Claridae	<i>Clariasbatrachus</i>	3.4	C	LC
36	Siluriformes/ Heteropneustidae	<i>Heteropneustesfossilis</i>	3.6	C	LC
37	Siluriformes/ Siluridae	<i>Ompokbimaculatus</i>	3.9	C	NT
38	Siluriformes/ Siluridae	<i>Wallagoattu</i>	3.7	C	NT
39	Siluriformes/ Schibeidae	<i>Eutropiichthysvacha</i>	3.9	C	LC
40*	Anabantiformes /Anabantidae	<i>Anabas testudineus</i>	3.0	M	LC
41	Cichliformes / Cichlidae	<i>Oreochromismossambicus</i>	2.2	M	NT
42	Cichliformes/ Cichlidae	<i>Pseudotropheus maculatus</i>	2.7	C	LC
43	Cichliformes/ Cichlidae	<i>Etroplusuratensis</i>	2.9	C	LC
44	Synbranchiformes / Mastacembelidae	<i>Mastacembelusarmatus</i>	2.8	C	LC
45	Synbranchiformes / Mastacembelidae	<i>Macrognathuspancalus</i>	3.5	C	LC
46	Perciformes /Ambassidae	<i>Chandanama</i>	3.9	C	LC
47	Perciformes /Ambassidae	<i>Parambassisranga</i>	3.6	C	LC


A= Abundant (76-100%); C = Common (51-75%); M = Moderate (26-50%); R = Rare (1-25%) of the total catch.

EN- Endangered; VU- Vulnerable; LC- Least concern; DD- Data deficient; NE- Not evaluated, NT: Near threaten.

\*Exotic fishes No.s: 4, 5 and 37

**Table: 2.** Number and percent composition of families, genera and species of fishes under various orders

S.No	Orders	% of families in an order	% of genera in an order	% of species in an order
1	Anguilliformes	5.26	2.94	4.25
	Beloiniformes	10.52	5.88	4.25
2	Cypriniformies	15.78	38.23	40.42
3	Cyprinodontiformes	5.26	2.94	2.12
4	Channiformes	5.26	2.94	6.38
5	Gobiiformies	5.26	2.94	2.12
6	Osteoglossiformes	5.26	2.94	2.12
7	Siluriformes	26.31	17.64	21.73
8	Anabantiformes	5.26	2.94	2.12
9	Cichliformes	5.26	8.82	6.38
10	Synbranchiformes	5.26	5.88	4.25
11	Perciformes	5.26	5.88	4.25




**Table: 3.** Taxonomic trophic level of fish species at Gosthani River

Trophic level	Herbivorous (2.0–2.5)	Omnivorous (2.6–3.5)	Carnivorous (3.6–4.50)
Number of species	09	22	16
% Composition	19.14	46.81	34.04

**Table: 4.** Taxonomic composition of population status and IUCN (2024)

Population Status	Abundant (76-100%)	Common (51-75%)	Moderate (26-50%)	Rare (1-25%)
Number of species	11	25	4	7
% Composition	23.40	53.19	8.51	14.89
IUCN (2024)	NT	LC	DD	NE
No. of species	06	39	01	01
% contribution	12.76	82.97	2.12	2.12



#### Ethical approval

This study was conducted according to international ethical standards set by the Institutional Plant care and Use Committee

#### Consent to participate

Not applicable as commercial gear operating reservoir, the local men and women were involved in the sampling study.

#### Data availability statement

The authors confirm that the data used to support the findings of this study are available within the manuscript.

**Comment [U5]:** Include a section on summary/conclusion to briefly give application benefits of the study to the fishery community surrounding the river.

#### REFERENCES

**Comment [U6]:** Put on fresh page

Burton PJ, AE Balisky, LP Coward, SG Cumming, DD Kneschwaw. The value of managing biodiversity. The Forestry Chronicle 1992; 68(2): 225-237.

CIFE. Assessment of riverine fisheries and linking with water quality restoration programme - river godavari in maharashtra. Final report. 2011: 1-110.

Chinnababu Sanapala., Dr Kantipriya Kondala., Dr Viswanadham Allada., Paidi Sandeep. Fish Diversity of Godavari River at Rajamahendravaram, Andhra Pradesh, India. 2021; 25 (6) : 21017 – 21022.

Chandra Sekhara Rao, Ch. Sebastian Raju and G. Simhachala. Biodiversity and Conservation Status of Fishes of River Sarada, Visakhapatnam District, Andhra Pradesh, India. Research Journal of Animal, Veterinary and Fishery Sciences, 2014; 2(2): 1-8.

Day, F., The fishes of India, being a natural history of the fishes known to inhabit the seas and freshwater of India, Burma and Ceylon, text and atlas, London, William Dawson and Sons Ltd., 1958: 195-198.

Ehrlich PR., and Wilson EO. Biodiversity studies science and policy. Science 1991; 253: 758-762.

Froese, R., Pauly, D., (eds.) 2020). Fish Base. [version 12/2020] <http://www.fishbase.org>  
<https://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>  
[https://www.alyoung.com/labs/biodiversity\\_calculator.html](https://www.alyoung.com/labs/biodiversity_calculator.html)

IUCN Red List of threatened species, version 2024. [www.iucnredlist.org](http://www.iucnredlist.org).

Jayaram, K. C. The freshwater fishes of the Indian region. Narendra Publication New Delhi, India, 1999: 551.

Jayaram K.C. The Freshwater Fishes of Indian Region Narendra Publication House, New Delhi, 2nd Edition, 2011.

Kaur, H., Datta, S.N. and Singh, A. Fish Catch Composition and Biodiversity Indices at Harike Wetland- A Ramsar Site in India. J. Anim. Res. 2017; 7(5): 935-941.

Menon A.G.K. Check list - freshwater fishes of India, Records of the Zoological Survey of India, Occasional, 1999:175- 366.

Mittermeier RA, CG Mittermeier., Megadiversity Earth's biologically wealthiest Nation. In McAllister, D.E. A. Lottamilion and B. Harvery (Eds)., Global S.V. Rankhamb /Rec Res Sci Tech 3 (2011) 11-1313 fresh water Biodiversity sea wind Cemex, Mexico city," 1997: 1-140.

Munro, I. S. R. The Marine and Freshwater Fishes of Ceylon. Biotech Books, Delh 2000.

Rama Rao, K., A study on fishing craft and gear in Lower Manair Dam, Karimnagar Dt. Andhra Pradesh, India. Int. J. M. Dis. Edu. Res. 2014 (a); 3: 5 (2): 56-68.

Rama Rao. K. Ichthyofaunal bio diversity in the lower Manair Dam at Karimnagar district; Telangana State: India Pelagia Research Library. Advances in Applied Science Research, 2014b; 5(5): 237-248.

Rama Rao K. Ichthyofaunal diversity in the Kalinga Dal reservoir at foothills of Mahendragiri, Mandasamandal, Srikakulam district, Andhra Pradesh State: India. International Journal of Fauna and Biological Studies 2018; 5(2): 89-96.

Rama Rao K. Ichthyofaunal diversity of Gotta Barrage at Hiramandalam, Vamsadhara River, Srikakulam Dt. Andhra Pradesh, India. International Journal of Zoology Studies, 2023; 8 (2): No. 23-28.

Rama Rao, K and R. Ramachandra Rao. Ichthyofaunal Diversity of Narayanapuram Anicut at Nagavali River, Srikakulam District of Andhra Pradesh, India. Uttar Pradesh Journal of Zoology, 2021;42(19): 24-35.

Rama Rao. K and Vinod Kumar Mudumala.. Ichthyofaunal biodiversity in the Sunamudi Gedda tributary of Mahendranaya, Eastren Ghats at Venkatavarada Rajapuram, Mandasa, Srikakulam Dt. Andhra Pradesh, India. International Journal of Multidisciplinary Educational Research, 2017;8(2):35-61.

Talwar P.K. and Jhingran A.G. Inland fishes of India and Adjacent Countries, Balkema, Rotterdam, Vol. A. 1991.

UNDER PEER REVIEW