

## Seasonal variation, diversity, and composition of planktons in the Ganga River

**Abstract:** Plankton are important components of aquatic ecosystems because they contribute to primary production, which sustains fisheries and other ecosystem functions. The study was carried out over a one-year period at three Ganga River sites: Haridwar (Bhadraabad) site A1, Bijnor (Balawali) site A2, and Muzaffarnagar (Bairaj Ganga bridge) site. Bacillariophyceae made almost 65% of the phytoplankton at location A1 14% Chlorophyceae and 14% Cynophyceae. The zooplankton consisted of Protozoa (80%) and Cladocera (20%), with Ulvophyceae accounting for 7%. At location A2, the phytoplankton composition was Bacillariophyceae (79%) and Chlorophyceae (7%). > Cynophyceae (7%). Ulvophyceae (7%), and zooplankton were Protozoa (80%), Rotifera (20%). At location A3, the phytoplankton mix shifted to Bacillariophyceae (70%). > Chlorophyceae (18%) > Cynophyceae (12%) and Protozoa (60%) Rotifera accounts for 40%. The Simpson diversity index (D) value for phytoplankton is 0.58, whereas for zooplankton, the value is 0.53. The mean plankton density at locations A1, A2, and A3 was 2059, 2959, and 3304 individuals per liter, respectively. Only dissolved oxygen in physicochemical characteristics correlated positively with plankton density; all other metrics correlated negatively.

**Keywords:** Simpson diversity, freshwater, density, zooplankton, phytoplankton

### Introduction

The most challenging task in ecology has been understanding the mechanisms that induce changes in biological communities' species richness and composition. Planktonic algae, the primary producers, are located at the bottom of the trophic pyramid while zooplankton were at higher layers of the pyramid and higher aquatic animals such as finfish, shellfish at the top. In addition to its biological variety, phytoplankton play a significant role in aquatic ecosystems by contributing to primary production, which supports fisheries and other vital features of ecosystems [1]. A rising number of studies are being conducted on developing large-scale river plankton [2, 3]. Compared to phytoplankton, zooplankton found in rivers receives less scientific attention. The stability of zooplankton in any aquatic body is profoundly important because they represent an important and sometimes unique food source for fish and many aquatic vertebrates [4]. Large, slow flowing rivers

and still bodies of water are frequently the only locations where well-developed phytoplankton communities may be found [5].

In contrast, higher flow rates can upset the phytoplankton structure. Plankton variety responds swiftly to changes in the aquatic environment, generally changes in water quality parameters, and reacts rapidly to pollution. Several planktonic species have been bioindicators [6, 7]. They are also essential for the biogeochemical cycles of several vital elements, including methanogenesis, nitrification, and the carbon cycle, and these cycles lead to primary production and recycling. since the biological communities can be considered as the representative of the environmental conditions, biological evaluation is a good indicator of the ecological quality of aquatic systems. Several recent studies on physico-chemical parameters and plankton community of rivers have been conducted on the Yamuna River [8], the Ganga River and its tributaries [9], the Sutlej River [10], and Jhelum River [11]. This study aimed to elucidate the changes in the composition of phytoplankton and zooplankton, their density, and their relationships with different physico-chemical parameters.

## Materials and Methods

This study was conducted for a twelve months period, from March 2019 to February 2020. Three sampling sites were selected in both rivers. In the Ganga river, three sites from each district were selected: *viz*: Haridwar (Bhadrabad) site A1, Bijnor (Balawali) site A2, and Muzaffarnagar (Bairaj Ganga bridge) site A3.

List 1 : Geographical locations of the sampling sites

Ganga river site	A1	A2	A3
Locations name	Bhadrabad	Balawali	Bairaj Ganga bridge
Map Location	29°55'15.1"N 78°04'42.2"E	29°38'07.0"N 78°06'21.7"E	29°22'26.1"N 78°02'03.5"E

Water samples were taken monthly from each sampling station to investigate the water quality parameters. Following standard procedure, we measured the sample site's electrical conductivity, pH, dissolved oxygen, free CO<sub>2</sub>, and water temperature [12]. Plankton samples were collected by filtering 100 L of the river water (through a plankton net of bolting silk number 25). The sieved

samples were transferred to the sampling bottles of 20 ml and preserved in 4% formalin and lugol's solution to ensure absolute preservation [12]. Phytoplankton was identified under 10X and zooplankton under 40X objective lenses of the compound microscope (Olympus- U-TV0.5XC-3). Plankton were counted with the help of a Sedgewick Rafter counting cell of 1 ml capacity. Identification and categorization of phytoplankton and zooplankton was done using the standard taxonomic keys of [13,14,15]. Karl Pearson's correlation coefficient and other statistical observations was measured using Microsoft Excel 2007 to determine the relationship among the various physico-chemical attributes.

Simpson's Diversity Index (D) measures diversity by considering the number of species present and each species' relative abundance.

The formula to calculate Simpson's Diversity Index (D) is

$$D = 1 - \left[ \frac{\sum n(n-1)}{N(N-1)} \right]$$

Where

n = the total number of organisms of a particular species

N = the total number of organisms of all species

## Results and Discussion

Monitoring the physico-chemical parameters to investigate how they affect the distribution of different biological components in freshwater ecosystems is crucial. The observed physico-chemical parameters from three different locations were DO (mg/l) recorded lowest at 7.4 and highest at 10.4, free carbon dioxide (mg/l) varied between 1.3 to 4.5, Water temperature (°C) between 13.1 to 20.1, pH between 7.2 to 7.9, and electrical conductivity (µS/cm) ranged between 122 to 189 during the entire study period. Higher dissolved oxygen values increase the photosynthetic rate of plankton and, hence, its density [16,17]. Phytoplankton use carbon dioxide and sunlight to make their food. The temperature range is significant because it impacts various metabolic processes in aquatic species [18]. Additionally, it safeguards aquatic life and modifies a stream's ecological health [19]. Alkaline water promotes high primary productivity [20]. EC is a

measure of the ability of water to pass electrical flow. Higher levels of total dissolved solids can often indicate pollution by an extraneous source [21]. [22] have reported the influence of various ranges of hydrobiological parameters on the density and composition of plankton. Monthly variation in plankton density and mean density of the Ganga River are presented in Table 1.

In Ganga river at site A1 14 species of Phytoplankton included Chlorophyceae (*Spirogyra sp.* and *Zygnema sp.*), Cynophyceae (*Phormedium sp.* and *Oscillatoria sp.*), Bacillariophyceae (*Asterionella sp.*, *Navicula sp.*, *Caloneis sp.*, *Cymbella sp.*, *Diatoma sp.*, *Stauroneis sp.*, *Nitzschia sp.*, *Fragilaria sp.*, *Hantzschia sp.*) and Ulvophyceae (*Cladophora sp.*) were found during the study period. The presence and absence of phytoplankton and zooplankton are shown in Tables 2 & 3, respectively. The highest abundance of Bacillariophyceae and a minimum of Ulvophyceae was observed at site A1. The order of phytoplankton occurrence was Bacillariophyceae (65%) > Chlorophyceae (14%) > Cynophyceae (14%) > Ulvophyceae (7%). At site A1, two zooplankton groups with five species were observed during the study period: Protozoa (80%) (*Astasia sp.*, *Colpoda sp.*, *Trachelophyllum sp.*, *Loxodes sp.*) and Cladocera (20%) (*Daphnia sp.*). Protozoa were the dominant group found at site A1. The mean concentration at site A1 was 2059 Individual/L. Plankton production was positively correlated with DO and negatively correlated with free CO<sub>2</sub>, Temperature, pH, and Electrical conductivity.

At site A2, 14 species of Phytoplankton included Chlorophyceae (*Spirogyra sp.*), Cynophyceae (*Phormedium sp.*), Bacillariophyceae (*Cocconeis sp.*, *Navicula sp.*, *Caloneis sp.*, *Cymbella sp.*, *Diatoma sp.*, *Stauroneis sp.*, *Nitzschia sp.*, *Fragilaria sp.*, *Diploneis sp.*, *Synedra sp.*, *Achnanthes sp.*) and Ulvophyceae (*Cladophora sp.*) were found during the study period. The presence and absence of phytoplankton and zooplankton are shown in Tables 4 & 5, respectively. At site A2 the highest abundance of Bacillariophyceae was observed and minimum abundance of Ulvophyceae. The order of occurrence of phytoplankton was Bacillariophyceae (79%) > Chlorophyceae (7%) > Cynophyceae (7%) > Ulvophyceae (7%). At site A2, 2 zooplankton groups with five species were observed during the study period: Protozoa (80%) (*Amoeba sp.*, *Colpoda sp.*, *Actinophrys sp.* and *Loxodes sp.*) and Rotifera (20%) (*Trichotria sp.*). Protozoa were the dominant group found at site A2. The mean density at site A1 was 2959 Individual/L. Plankton production positively correlated with DO and negatively correlated with free CO<sub>2</sub>, Temperature, pH, and Electrical conductivity.

At site A3, 17 species of Phytoplankton included Chlorophyceae (*Spirogyra sp.*, *Closteriopsis sp.* and *Pediastrum sp.*), Cynophyceae (*Phormedium sp.* and *Spirulina sp.*) and Bacillariophyceae (*Cyclotella sp.*, *Frustulia sp.*, *Navicula sp.*, *Caloneis sp.*, *Cymbella sp.*, *Diatoma sp.*, *Stauroneis sp.*, *Nitzschia sp.*, *Fragilaria sp.*, *Achnanthes sp.*, *Cocconeis sp.*, and *Diatomella sp.*) were found during the study period. The presence and absence of phytoplankton and zooplankton are shown in Tables 6 & 7, respectively. At site A3, the Bacillariophyceae group was most abundant, whereas the Cynophyceae group was least abundant. The order of phytoplankton occurrence was Bacillariophyceae (70%) > Chlorophyceae (18%) > Cynophyceae (12%). At site A3, 2 zooplankton groups with six species were observed during the study period: Protozoa (60%) (*Amoeba sp.*, *Colpoda sp.*, *Astasia sp.* and *Trachelophyllum sp.*) and Rotifera (40%) (*Dicranophous sp.* and *Testudinella sp.*). Protozoa were the dominant group found at site A2. The mean density at site A3 was 3304 Individuals/L. We observed plankton production, which was found to have a positive correlation with DO and a negative correlation with free CO<sub>2</sub>, temperature, pH, and electrical conductivity.

The Simpson diversity index (D) value for phytoplankton at site A1 (D= 0.58), A2 (0.40), and A3 (0.49) shows that the highest diversity of phytoplankton was at site A1 and lowest at site A2. For zooplankton, the value of D at site A1 (D= 0.40), A2 (0.40), and A3 (0.53) shows that the highest diversity for zooplankton was observed at site A1 and lowest at site A3.

The highest density in the Ganga River was observed during January and the lowest during the monsoon season, particularly in September, as presented in Table 1; similar results were obtained by [23,24]. In September, similar observations was reported from the Madurai water body [25]. This increase is due to favorable conditions returning after the monsoon season. The highest abundance of Bacillariophyceae has been observed at sites A1, A2, and A3; this is because, in contrast to other groups, Bacillariophyceae can adapt to a wide range of environmental conditions. [26] claim that the Bacillariophyceae class is cosmopolitan due to its excellent tolerance, adaptability, and ecological adaptation. [9] reported a total absence of rotifers during the monsoon season in the Ganga River, as seen in the present study at site A2. [27,28,29] among others, have also reported a similar trend. The maximum density of 4844 Individuals/L in Ganga River at the site in January, a similar trend was observed by [30]. In the present study, phytoplankton was found to be dominant over zooplankton. According to [31] zooplankton is one



1. <i>Spirogyra</i> sp.	++	+	++	++	+	+	-	+	+	++	++	+
2. <i>Zygnema</i> sp.	-	-	+	+	+	-	-	-	+	++	++	-
<b>Cynophyceae</b>												
1. <i>Phormedium</i> sp.	++	+	++	++	+	-	-	-	-	+	++	++
2. <i>Oscillatoria</i> sp.	+	+	+	+	+	-	+	+	-	-	-	+
<b>Bacillariophyceae</b>												
1. <i>Asterionella</i> sp.	++	++	+	+	+	-	+	-	+	+	+	-
2. <i>Navicula</i> sp.	+	+	+	-	+	+	-	+	-	-	+	++
3. <i>Caloneis</i> sp.	++	+	++	++	+	+	+	+	++	++	++	+
4. <i>Cymbella</i> sp.	+	+	++	++	+	+	-	+	+	+	++	++
5. <i>Diatoma</i> sp.	+	++	++	++	+	+	+	+	+	+	++	++
6. <i>Stauroneis</i> sp.	+	+	-	-	-	-	-	-	-	+	+	+
7. <i>Nitzschia</i> sp.	+	+	++	++	+	+	+	+	++	++	++	++
8. <i>Fragilaria</i> sp.	++	+	+	+	+	-	-	-	-	-	++	+
9. <i>Hantzschia</i> sp.	+	++	++	++	+	+	+	-	-	+	+	-
<b>Ulvophyceae</b>												
1. <i>Cladophora</i> sp.	+	++	++	+	-	-	-	+	+	-	+	-

Note: - (Absent)  
+ (Present)  
++ (Abundant)

**Table 3. Zooplankton composition observed at site A1 of the Ganga River**

	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb
<b>Cladocera</b>												
1. <i>Daphnia</i> sp.	+	-	+	+	-	-	-	-	+	-	+	+
<b>Protozoa</b>												
1. <i>Astasia</i> sp.	++	+	++	+	++	+	-	-	-	+	++	++
2. <i>Colpoda</i> sp.	+	+	++	+	+	-	+	+	+	++	++	+
3. <i>Trachelophyllum</i> sp.	+	+	-	-	-	-	-	-	-	-	+	+
4. <i>Loxodes</i> sp.	-	-	--	++	+	+	-	-	-	-	-	+

Note: - (Absence)  
+ (Presence)  
++ (Abundance)

**Table 4. Phytoplankton composition observed at site A2 of the Ganga River**

	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb
<b>Chlorophyceae</b>												
1. <i>Spirogyra sp.</i>	++	++	+	++	+	+	-	+	+	+	++	++
<b>Cynophyceae</b>												
1. <i>Phormidium sp.</i>	+	+	+	+	+	-	-	-	-	+	+	++
<b>Bacillariophyceae</b>												
1. <i>Cocconeis sp.</i>	+	+	++	+	+	-	+	-	+	+	+	-
2. <i>Navicula sp.</i>	++	+	+	-	+	+	-	+	-	-	+	+
3. <i>Caloneis sp.</i>	++	+	+	+	+	-	-	+	+	-	+	++
4. <i>Cymbella sp.</i>	+	+	++	++	+	+	+	+	+	++	++	++
5. <i>Diatoma sp.</i>	++	+	+	++	++	+	+	+	++	+	+	+
6. <i>Stauroneis sp.</i>	+	+	+	+	+	+	+	+	+	-	-	+
7. <i>Nitzschia sp.</i>	-	+	+	-	-	-	-	+	++	++	+	-
8. <i>Fragilaria sp.</i>	+	+	+	-	-	-	-	-	-	++	+	++
9. <i>Diploneis sp.</i>	++	++	+	+	+	-	-	-	+	++	++	++
10. <i>Synedra sp.</i>	+	+	+	+	+	+	+	+	+	+	+	+
11. <i>Achnanthes sp.</i>	+	+	-	-	-	-	-	+	+	++	+	-
<b>Ulvophyceae</b>												
1. <i>Cladophora sp.</i>	+	+	+	+	-	-	-	+	+	-	+	-

Note: - (Absence)  
 + (Presence)  
 ++ (Abundance)

**Table 5. Zooplankton composition observed at site A2 of the Ganga River**

	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb
<b>Rotifera</b>												
1. <i>Trichotria sp.</i>	+	-	+	+	-	-	-	-	+	-	++	+
<b>Protozoa</b>												
1. <i>Amoeba sp.</i>	+	++	+	++	+	+	-	-	-	+	++	+
2. <i>Colpoda sp.</i>	-	+	++	+	-	-	+	+	++	++	++	+
3. <i>Actinophrys sp.</i>	+	+	-	-	-	+	+	+	+	-	-	+
4. <i>Loxodes sp.</i>	-	-	-	+	+	-	-	-	-	-	+	+

Note: - (Absence)  
 + (Presence)  
 ++ (Abundance)

**Table 6. Phytoplankton composition observed at site A3 of the Ganga River**


	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb
<b>Chlorophyceae</b>												
1. <i>Spirogyra sp.</i>	+	++	++	++	+	+	-	+	++	++	+	++
2. <i>Closteriopsis sp.</i>	+	+	-	-	-	+	+	+	+	-	-	-
3. <i>Pediastrum sp.</i>	+	+	+	+	-	-	-	-	-	-	++	+
<b>Cynophyceae</b>												
1. <i>Phormedium sp.</i>	+	+	+	+	+	-	-	-	-	+	+	+
2. <i>Spirulina sp.</i>	-	+	++	++	+	-	-	-	-	+	+	+
<b>Bacillariophyceae</b>												
1. <i>Cyclotella sp.</i>	+	+	+	+	+	-	+	-	+	++	++	-
2. <i>Frustulia sp.</i>	+	+	+	-	+	+	-	+	-	-	+	+
3. <i>Caloneis sp.</i>	+	+	+	+	+	-	-	+	++	+	++	++
4. <i>Cymbella sp.</i>	++	++	+	++	+	+	+	+	+	+	+	+
5. <i>Diatoma sp.</i>	+	++	+	+	+	+	+	+	+	++	-	+
6. <i>Fragilaria sp.</i>	+	++	++	+	+	+	+	+	+	++	++	++
7. <i>Achnanthes sp.</i>	+	+	+	+	+	-	-	-	-	+	+	+
8. <i>Navicula sp.</i>	-	++	+	++	+	-	-	-	-	+	++	++
9. <i>Nitzschia sp.</i>	+	+	+	+	+	+	+	+	+	-	-	+
10. <i>Stauroneis sp.</i>	++	+	++	++	+	-	-	-	+	+	++	++
11. <i>Cocconeis sp.</i>	+	+	++	+	++	+	-	-	-	+	++	-
12. <i>Diatomella sp.</i>	-	-	+	+	+	-	-	-	+	+	-	+

Note: - (Absence)  
 + (Presence)  
 ++ (Abundance)

**Table 7. Zooplankton composition observed at site A3 of the Ganga River**

	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb
<b>Rotifera</b>												
1. <i>Dicranophous sp.</i>	+	+	-	-	-	-	-	-	+	++	++	-
2. <i>Testudinella sp.</i>	-	+	+	-	-	+	+	-	-	-	-	+
<b>Protozoa</b>												
1. <i>Amoeba sp.</i>	+	+	+	+	+	+	-	-	-	+	+	+
2. <i>Colpoda sp.</i>	-	+	+	+	-	-	+	+	+	++	++	+
3. <i>Astasia sp.</i>	+	-	+	+	+	-	-	-	++	+	++	+
4. <i>Trachelophyllum sp.</i>	+	++	+	+	+	-	-	-	-	-	-	-

Note: - (Absence)  
 + (Presence)  
 ++ (Abundance)



*Achnanthes sp.*




*Actinophrys sp.*



*Amoeba sp.*



*Astartia sp.*




*Asterionella sp.*



*Caloneis sp.*



*Cladophora sp.*



*Closteriopsis sp.*



*Coelastrum sp.*




*Colpoda sp.*



*Cyclotella sp.*



*Cymbella sp.*



*Diatoma sp.*




*Diatomella sp.*



*Dicrannophorus sp.*



*Diploneis sp.*



*Fragilaria sp.*



*Frustulia sp.*



*Hantzchia sp.*




*Loxodes sp.*




*Navicula sp.*



*Nitzschia sp.*



*Oscillatoria sp.*



*Pediastrum sp.*



*Phormidium sp.*



*Spirulina sp.*



*Stauroneis sp.*



*Synedra sp.*



*Trachelophyllum sp.*




*Trichotria sp.*




*Spirogyra sp.*



*Zygnuma sp.*



*Cocconeis sp.*



*Testudinella sp.*



*Daphnia sp.*

**Figure 1** Planktons of Ganga river found during the study period

### Conclusion

In the present study, Bacillariophyceae emerged as the dominant group of phytoplankton, and protozoa were the dominant group among zooplankton among the studied stations. The mean plankton density at sites A1, A2, and A3 were 2059, 2959, and 3304 individuals/L respectively, showing site A3 favors plankton growth. Simpson diversity index (D) value for phytoplankton shows the highest diversity at site A1 (D= 0.58). It shows that phytoplankton of a wide variety can flourish here, and the same has been observed for zooplankton at site A3 (0.53). It shows phytoplankton and zooplankton of a wide variety can flourish at each site. The basic information on the phytoplankton and zooplankton distribution and abundance would be helpful for further ecological assessment and monitoring of freshwater ecosystems.

### References

1. Dhargalkar VK, Verlecar XN. Zooplankton Methodology, Collection & Identification-a field manual. National Institute of Oceanography, Dona Paula, Goa. 2004.
2. Basu BK, Kalff J, Pinel-Alloul B. Midsummer plankton development along a large temperate river: the St. Lawrence River. Canadian Journal of Fisheries and Aquatic Sciences. 2000. 14;57(S1):7-15.
3. Reckendorfer, W., Keckeis, H., Winkler, G. and Schiemer, F. Abundance in the River Danube, Austria: Ce of inshore retention. Freshw. Biol. 1999. 41, pp.583-591.

4. Ochang, S.N., Ayotunde, E.O. and Okey, I.B. Some aspect of the physico-chemical and biological properties of Cross river, an inland water body in South-eastern Nigeria. *Global Journal of Agricultural Sciences*. 2005. 4(2), pp.139-148.
5. Wetzell, R. G. *Limnology: Lake and river Ecosystem*, 3rd ed. Academic Press. 2001. ISBN -12-744760-1.
6. Vareethiah, K. and Haniffa, M.A. Phytoplankton community organisation and species succession in a Bar-built estuary. *Journal of Environment and Pollution*. 1998. 5, pp.209-214.
7. Bianchi, F., Acri, F., Aubry, F.B., Berton, A., Boldrin, A., Camatti, E., Cassin, D. and Comaschi, A. Can plankton communities be considered as bio-indicators of water quality in the Lagoon of Venice. *Marine Pollution Bulletin*. 2003. 46(8), pp.964-971.
8. Kumar, S. and Saxena, A. Plankton Composition in Relation to Physio-Chemical Parameters of Yamuna River at Three Different Cities of India. *Research Biotica*. 2022. 4(3), pp.94-101.
9. Khanna, D. R., Rakesh Bhutiani Rakesh Bhutiani, Gagan Matta Gagan Matta, Vikas Singh Vikas Singh, and Gaurav Bhadauriya Gaurav Bhadauriya. Study of planktonic diversity of river Ganga from Devprayag to Roorkee, Uttarakhand (India). 2012: 211-217.
10. Kaur, S. and Singh, P. Studies on plankton diversity of River Sutlej, Punjab. *Journal of Entomology and Zoology Studies*. 2017. 5(6), pp.620-628.
11. Rashid, H. and Pandit, A.K. Ecology of plankton community of river Sindh in Kashmir Himalaya. *J. Himalayan Ecol. Sustian*. 2008. Dev, 3, pp.11-22.
12. APHA. *Standard methods for the examination of the water and wastewater*. 22th edition. American Public Health Association, Washington Aquaculture Engineering. 2012. p. 19.
13. Edmondson, W.T. *Freshwater Biology*. John Wiley and Sons Inc., New York. 1959. 1248 p
14. Pennak R.W. *Freshwater Invertebrates of United State*. 2<sup>nd</sup> edition John Wiley & Sons, New York. 1978. 822pp.
15. Biggs, B.J. and Kilroy, C. *Stream periphyton monitoring manual*. Niwa. 2000.
16. Sharma, R.K. and Rathore, V. Pollution ecology with reference to commercially important fisheries prospect in a rural based water body: The Lake Sarsai Nawar, Etawah In U. P.(India). *Pollution Research*. 2000. 19(4), pp.641-644.

17. Ravindra, K., Mor, S., Ameena, Kamyotra, J.S. and Kaushik, C.P. Variation in spatial pattern of criteria air pollutants before and during initial rain of monsoon. *Environmental Monitoring and Assessment*. 2003. 87, pp.145-153.
18. Gupta, M., Kumar, S., Dangi, S.S. and Jangir, B.L. Physiological, biochemical and molecular responses to thermal stress in goats. *International Journal of Livestock Research*. 2013. 3(2), pp.27-38.
19. Hanjra, M.A., Blackwell, J., Carr, G., Zhang, F. and Jackson, T.M. Wastewater irrigation and environmental health: Implications for water governance and public policy. *International journal of hygiene and environmental health*. 2012. 215(3), pp.255-269.
20. Kumar, M.P. and Prabhakar, C. Physico-chemical parameters of river water: a review. *Int J Pharm Biol Arch*. 2012. 3, pp.1304-1312.
21. Aravinth, A., Kannan, R., Chinnadurai, G., Manickam, N., Raju, P., Perumal, P. and Santhanam, P. Temporal changes in plankton diversity in relation to hydrographical characteristics at Perumal Lake, Cuddalore District, Tamil Nadu, India. *The Journal of Basic and Applied Zoology*. 2023. 84(1), p.13.
22. Anil, P., Madhu, N.V., Vishal, C.R., Gopika, P., Jyothi, S., Arya, K.S. and Gireeshkumar, T.R. Characterization of phytoplankton functional groups in a tropical shellfish harvesting estuary (Ashtamudi) and adjacent nearshore waters (southwest coast of India). *Environmental Science and Pollution Research*. 2023. 30(12), pp.34553-34572.
23. Fitt, W.K., McFarland, F.K., Warner, M.E. and Chilcoat, G.C. Seasonal patterns of tissue biomass and densities of symbiotic dinoflagellates in reef corals and relation to coral bleaching. *Limnology and oceanography*. 2000. 45(3), pp.677-685.
24. Yogendra, K. and Puttaiah, E.T. Determination of water quality index and suitability of an urban waterbody in Shimoga Town, Karnataka. In *Proceedings of Taal 2007: The 12th world lake conference*. 2008. (Vol. 342, p. 346).
25. Sharma, S., Siddique, A., Singh, K., Chouhan, M., Vyas, A., Solnki, C.M., Sharma, D., Nair, S. and Sengupta, T. Population dynamics and seasonal abundance of zooplankton community in Narmada River (India). *Researcher*. 2010. 2(9), pp.1-9.

26. Arinardi, O.H., Sutomo, A.B., Yusuf, S.A., Trimaningsih, E.A. and Riyono, S.H. The range of abundance and predominant plankton composition in the waters of Eastern Indonesia. Jakarta: P3O-LIPI. 1997.
27. Welch, E.B. Phytoplankton and related water-quality conditions in an enriched estuary. *Journal (Water Pollution Control Federation)*. 1968. pp.1711-1727.
28. Srivastava, K., Mishra, S., Verma, H.O., Thakur, V.R., Jha, D.N., Alam, A. and Das, B.K. Time scale changes of plankton in the river Ganga at Kanpur. 2020.
29. Mohanta, K.N., Subramanian, S., Komarpant, N. and Saurabh, S. Alternate carp species for diversification in freshwater aquaculture in India. *Aquaculture Asia*. 2008. 13(1), p.11.
30. Magar, V.G. Study on The Relationship of Plankton diversity and Physico-Chemical Parameters of Barakune Daha, Dang District, Nepal (Doctoral dissertation, Department of Zoology). 2009.
31. Brysiewicz, A., Jankowski, M. and Tański, A. Zooplankton structure in midfield ponds in north-west part of Poland. *Infrastruktura i Ekologia Terenów Wiejskich*. 2017. (IV/2), pp.1673-1686.
32. Anyinkeng, N., Mih, A.M., Suh, T.A. and Awah, C.C. Phytoplankton diversity and abundance in water bodies as affected by anthropogenic activities within the Buea municipality, Cameroon. *Journal of Ecology and the Natural Environment*. 2016. 8(7), pp.99-114.
33. Pandit, D.N., Kumari, P. and Sharma, S.K. Ecology and diversity of Zooplankton of the river Ganga at Bihar, India in Relation to Water Quality. *Current World Environment*. 2020. 15(2), p.304.
34. Kudari, V.A., Kadadevaru, G.G. and Kanamadi, R.D. Zooplankton composition in some ponds of Haveri district, Karnataka. *Zoo's print Journal*. 2005. 20(12), pp.2094-2099.
35. Baruah, P.P., Kalita, H. and Nath, S. A taxonomic account on phytoplankton of Rudrasagar Ramsar site, Tripura (India): Chlorophyta and Euglenophyta. *The Journal of Indian Botanical Society*. 2020. 100(3and4), pp.91-118.
36. Chowdhury, M.M.R., Mondol, M.R.K. and Dewan, S. Seasonal dynamics of plankton in relation to some environmental factors in a Beel ecosystem. *University Journal of Zoology, Rajshahi University*. 2008. 27, pp.55-58.

37. Sharma C, Jindal R, Singh UB, Ahluwalia AS, Thakur RK. Population dynamics and species diversity of plankton in relation to hydrobiological characteristics of river Sutlej, Punjab, India. *Ecol Environ Conserv*. 2013. 19(3):717-24.
38. Elmoor-Loureiro LM, Sousa FD, Oliveira FR, Joko CY, Perbiche-Neves G, Da Silva AC, Silva AJ, Ghidini AR, Meira BR, Aggio CE, Morais-Junior CS. Towards a synthesis of the biodiversity of freshwater Protozoa, Rotifera, Cladocera, and Copepoda in Brazil. *Limnologica*. 2023 May 1;100: 126008.

UNDER PEER REVIEW