

A Comprehensive Review on Scrub Typhus: Epidemiology, Transmission, Diagnosis and Management

Abstract:

Scrub typhus is a serious public health risk in areas where it is endemic. It is brought on by the bacterium *Orientia tsutsugamushi* and is spread via the bite of infected chiggers. The epidemiology, clinical symptoms, diagnosis, treatment, and prevention of scrub typhus are summarized in this overview, along with recent developments. The disease is most common in rural sections of Asia, the Pacific Islands, and portions of Australia, where chigger populations are encouraged by favorable environmental conditions.

The clinical signs of chigger bites can range in severity from mild to severe, presenting with symptoms like fever, headaches, and the distinctive Escher at the bite site. Severe cases may result in consequences such as acute respiratory distress syndrome and organ failure, which call for prompt identification and treatment. Molecular techniques and serological testing are used as diagnostic tools. Molecular assays are a quick and reliable way to identify *Orientia tsutsugamushi*.

Antibiotics are the cornerstone of treatment, and doxycycline is the drug of choice. Treatment must be started as soon as possible to avoid problems and enhance results. The importance of public health interventions, such as vector control and health education, is highlighted by issues like antibiotic resistance and restricted access to treatment in rural areas.

Personal protection is the main focus of preventive measures, which include donning proper attire and applying insect repellents. Managing the environment and involving the community are important strategies for lowering chigger populations. Additional research is being done to create effective scrub typhus vaccines, even if there isn't one that is now generally authorized.

Key words:

Orientia tsutsugamushi, Scrub Typhus, Escher, Vector-Borne Disease, and Chigger-Borne Disease, rickettsial infection, zoonosis, diagnostic tests, scrub typhus epidemiology, endemic regions, and vaccine development.

1. Introduction:

Orientia tsutsugamushi, a member of the Rickettsiaceae family of bacteria, is the infecting agent that causes scrub typhus. Through the bite of an infected chigger, a mite in the larval stage, people can contract this zoonotic disease(1). *Scrub typhus is the name given to these mites* because they are most common in high-humidity environments like grasslands, dense flora, and scrubby terrain. "*Orientia tsutsugamushi*, the bacteria that causes scrub typhus, is an obligatory intracellular infection(2). Its life cycle is distinct in that it involves rodents as reservoir hosts and trombiculid mites, or chiggers, as vectors for transferring the bacteria to humans(3). The bacteria mainly affect the host's endothelium and mononuclear cells, which might result in a variety of clinical symptoms. Scrub typhus is primarily found in the Asia-Pacific region, which includes countries such as China, India,

Japan, and some Southeast Asian countries. Although cases have been reported in the Pacific Islands and some regions of the Middle East, these are the places where it is most common. Scrub typhus is mostly dependent on environmental variables; the mites thrive in regions with high humidity and dense vegetation(4).

The clinical manifestations of scrub typhus usually appear six to twenty-one days following the bite of an infected chigger. Frequent signs and symptoms include chills, headaches, muscle aches, and abrupt fever onset. The emergence of an eschar, a black, scab-like lesion at the site of the chigger bite, is a hallmark that sets scrub typhus apart. This eschar helps to distinguish scrub typhus from other febrile diseases and is a crucial diagnostic signal. Scrub typhus has the potential to worsen and impact several organ systems if left untreated(5). Meningoencephalitis, acute renal damage, respiratory distress, and disseminated intravascular coagulation are among the possible complications. Scrub typhus can be fatal in extreme cases, which highlights the significance of early detection and timely treatment.

Diagnosing scrub typhus involves laboratory tests, with serological assays being commonly employed. These tests are used to identify antibodies that the immune system produces in reaction to the infection. The genetic makeup of the bacterium can also be identified by molecular methods like polymerase chain reaction (PCR)(6). Antibiotics are the main treatment for scrub typhus, and the best medication for this condition is doxycycline. Other antibiotics like azithromycin or chloramphenicol may be utilized when doxycycline treatment is contraindicated. The prevention of illness development and the mitigation of associated consequences necessitate the early beginning of antibiotic therapy. Scrub typhus can be avoided by combining environmental management with personal preventive measures. It is recommended that people stay away from densely vegetated regions, wear protective gear (long sleeves and pants), apply insect repellents, and thoroughly check themselves for chiggers after engaging in outdoor activities. Additionally, in an effort to lower the number of mites and stop the bacteria from spreading, public health initiatives concentrate on environmental cleaning and rodent control(7).

1.2. History and background:

In Asia, scrub typhus has long been known as a disease. Historical reports indicate that diseases like scrub typhus were prevalent in antiquity, particularly in parts of China and Japan. Scrub typhus was difficult to distinguish from other fever infections because these early reports frequently confused the disease.

In the 20th century, the phrase "scrub typhus" was first used. The disease gained considerable attention during World War II, when outbreaks occurred among soldiers stationed in the Asia-Pacific region. The term "scrub typhus" refers to the terrain—wooded, scrubby areas with a high chigger population—where the disease frequently manifests(8).

1.3. Discovery of the Disease Agent:

Japanese scientist H. Hayashi discovered *Orientia tsutsugamushi*, the causal agent of scrub typhus, in 1930. After separating the bacteria from the patients, he determined that the infection was causing the illness(9). This finding was a significant advancement in our knowledge of scrub typhus. Scrub typhus has continued to be a public health hazard in many Asian countries since it was discovered, particularly in rural and forested areas. Scrub typhus-related mortality has been dramatically decreased by early identification and effective antibiotic treatment(10).

1.4. Geographical distribution:

1.4.1. Japan:

Japan is the country where scrub typhus was first discovered, and there are still cases of the disease there. Due to an increase in mite populations transmitting *O. tsutsugamushi*'s, there was a notable comeback and outbreak between 1976 and 1984(11). The increased quantity of mites during that time period lacked an explanation. Almost all of Japan is currently affected by the disease, with the exception of the prefectures of Okinawa and Hokkaido. A 1998 retrospective analysis showed that the highest percentage of cases was found in Kyushu (51% of total cases), Tohoku-Hokuriku (27%), and Kanto (19%). The high number of cases in Kanto and Kyushu, in contrast to China, accounted for the biggest percentage of reported cases in November(12).

1.4. 2. Korea:

Reports of scrub typhus have been made in North and South Korea, with the majority of cases occurring in rural and mountainous areas. Korean civilians were not aware of scrub typhus until 1986, despite the fact that cases were first documented in South Korea during the Korean War(13). As a result, the illness is now acknowledged as the most prevalent rickettsia illness in South Korea. Between 1986 and 1993, *O. tsutsugamushi*'s was found to be seropositive in 27.7% to 51% of acute fever patients in South Korea, according to nationwide seroepidemiologic and microbiologic surveys. Scrub typhus is a widespread illness in the nation that is often under diagnosed, according to the research. South Korea started to report cases of scrub typhus in 1994(14). Physicians are required to notify the Korean Centres for Disease Control and Prevention (CDC) and the local health bureau of any confirmed or suspected cases of scrub typhus(15).

1.4.3. India:

In India, scrub typhus is a common disease in rural areas; occurrences have been documented in Himachal Pradesh, Uttarakhand, Jammu and Kashmir, and the north eastern provinces. In 1917, scrub typhus was identified as a typhus-like disease in India(16). During World War II and the Indo-Pak War of 1965, it was a significant cause of fever among military men along the border between Assam, India, Myanmar (formerly Burma), and Myanmar. In 1990, the disease reappeared at India's border with Pakistan(17). This decline in incidence was caused by alterations in lifestyle, the widespread use of insecticides, and the empirical treatment of febrile illness. Scrub typhus is still a disease that is not well recognized in India(18). The disease is found throughout India, from the south to the northeast and northwest, according to field epidemiology research. Maharashtra, Tamil Nadu, Karnataka, Kerala, Himachal Pradesh, Uttaranchal, Rajasthan, West Bengal, Bihar, Meghalaya, and Nagaland were among the states from which instances were reported(19). The months of August through October are when the illness peaks. It has been stated that *O. tsutsugamushi*'s major vector is *Leptotrophilium deliense*. Two significant risk factors for scrub typhus are occupation and socioeconomic level. The majority of Indian scrub typhus patients are illiterate and from rural areas(20).

1.4.4. Other countries:

The *tsutsugamushi*'s triangle includes a good number of additional nations with reported cases of scrub typhus in addition to those mentioned above. Scrub typhus has been known for about a century on the Australian continent as well as the islands of the southwest Pacific, such as Indonesia and the Philippines. In Australia, it was identified as "coastal fever" in 1913 and as scrub typhus after the

1920s. The tropical coastal regions of north eastern Queensland, the tropical Northern Territory, and the neighbouring Kimberly region of Western Australia are the endemic areas in Australia(21). In 1998, Litchfield, an island nation distinct from others in the Asia-Pacific region, was discovered in Australia. Scrub typhus was first identified in the Philippines during World War II. The lungs and spleens of white rats infected with the vainer strain of *O. tsutsugamushi*'s were used to create the first, unsuccessful US scrub typhus vaccine. In the Philippines, the vainer strain was first discovered in a soldier's blood. Scrub typhus in Malaysia has a history that dates back to 1915. In the Solomon Islands, Republic of Vanuatu, and Papua New Guinea, the sickness gained notoriety during World War II(22).

1.5. Causative agent:

Orientia tsutsugamushi is the bacterium that causes scrub typhus. Humans contract this bacterium when bitten by a bite from infected larvae of the trombiculid mite, commonly referred to as chiggers. For the most part, scrub typhus is restricted to the Asia-Pacific region, which includes northern Australia, sections of Southeast Asia, the Indian subcontinent, and several Pacific islands. The bacterium *Orientia tsutsugamushi* is an obligatory intracellular one, which means it can only live and grow inside host cells(23). Scrub typhus, the disease it produces, is typified by symptoms including fever, headache, soreness in the muscles, and a distinctive eschar, or black, scab-like lesion, at the site of the chigger bite. Scrub typhus can cause major consequences, including organ failure, if treatment is not received. Avoiding chigger-infested regions, donning protective clothes, and applying insect repellents are examples of preventive strategies. Antibiotics, such as doxycycline or azithromycin, are commonly used in treatment. These drugs are efficient in battling infections when given early in the course of the illness(24).

1.6. Life cycle:

Scrub typhus is caused by *Orientia tsutsugamushi*, a bacterium that has two hosts in its life cycle: an invertebrate host, which is the larval stage of trombiculid mites, also referred to as chiggers, and a vertebrate host, which is typically a rat(25). The life cycle and method of transmission of scrub typhus by chigger bites are briefly described below:

1.6.1. Chigger Larvae Feeding Stage:

The life cycle of trombiculid mites, sometimes known as chiggers, starts when the larval stage feeds on the blood of an infected vertebrate host, usually a small animal like a mouse. The bacterium, *Orientia tsutsugamushi*, is acquired by the larvae during this blood feeding.

1.6.2. Infection in Chiggers:

The germs that are consumed grow inside the chigger larvae, causing an illness. The mites develop into bacterium-carrying vectors after becoming infected.(26)

1.6.3. Nymph and Adult Stages of Chiggers:

The chigger larvae fall to the ground and molt into nymphs and adult mites after feasting on the blood of their vertebrate host.

1.7. Transmission to Vertebrate Hosts:

Infected chiggers look for a fresh vertebrate host to feed on, usually small animals like mice. They can transfer the bacterium to the vertebrate host while they are feeding(27).

1. 7. 1. Maintenance in Vertebrate Hosts:

The bacterium cause infection and has the ability to spread throughout the bloodstream in its vertebrate host. The natural reservoir hosts of *Orientiatsutsugamushi* are usually small animals.

1. 8. Transovarial Transmission:

The bacteria may occasionally be passed on to the following generation of mites through a process known as transovarial transmission, which occurs when contaminated female mites transfer the bacteria to their eggs.

1. 9. Human Infection:

While chiggers bite humans while they are still in the larval stage, they can contract the infection. During feeding, the chigger's saliva carries the infection.

1. 9. 1. Clinical Manifestation in Humans:

Orientiatsutsugamushi causes scrub typhus in humans, which manifests as fever, headache, muscle soreness, and the development of an eschar at the site of the chigger bite.

1. 10. Transmission through chigger bites

1. 10. 1. Chigger Attachment:

Because they are so small, larval chiggers often go unnoticed. They typically like to adhere to human skin in warm, moist areas like skin folds.(28)

1. 10. 2. Feeding Procedure:

The chigger makes a puncture in the skin and injects saliva, which contains many enzymes that break down the host's tissue. The chigger injects some of its saliva back into the skin of its host during feeding, in addition to sucking up the liquefied tissue.

1. 10. 3. Bacterial Transmission:

The bite wound allows the bacteria-containing saliva from the chigger to enter the human host's bloodstream if it is infected with *Orientiatsutsugamushi*.

1. 11. Infection and Disease:

The bacteria enter the body through the lymphatic and blood vessels' endothelial cells, infecting them and producing the symptoms of scrub typhus. Scrub typhus can be prevented by avoiding areas with a lot of grass and bushes, wearing protective clothing (long sleeves and pants), using insect repellents, and carefully checking and getting rid of any chiggers after visiting areas that may be infected. Anyone with symptoms such as fever, rash, or headache and suspecting they have been bitten by chiggers should seek medical attention immediately in order to receive a diagnosis and appropriate treatment, including antibiotics(29).

1.12. Vector Host

1. 12. 1. Vector Host (Chiggers):

1. 12. 2. Larval Stage:

The main vectors for scrub typhus transmission are chiggers. They consume the blood of vertebrate hosts, including humans, during their parasitic larval stage.

1. 12. 3. Infection:

When chiggers feed on the blood of an infected vertebrate host, usually a small mammal like a rodent, they contract *Orientiatsutsugamushi*.

1. 12. 4. Multiplication of Bacterium:

The bacterium grows inside the chigger and creates an infection that can spread to other chiggers when they feed again.

1.13. Transmission to Vertebrate Hosts:

1. 13. 1. Feeding Process:

Chiggers carrying the infection look for fresh vertebrate hosts, frequently small mammals, to consume. They inject saliva into the host—which might contain *Orientiatsutsugamushi*—during feeding.

1. 13. 2. Transmission:

The chigger's saliva carries the bacteria to its new vertebrate host. This causes infection in the vertebrate host, which includes rodents and other small mammals.

1. 13. 3. Vertebrate Hosts:

1. 13. 4. Natural Reservoir:

Orientiatsutsugamushi natural reservoir hosts are small mammals, especially rodents. Chiggers that feed on these animals can become infected due to the bacterium that circulates in their bloodstream.

1. 13. 5. Maintenance of Bacterium:

In endemic regions, scrub typhus persists because the bacterium is kept alive in the population of vertebrate hosts.

1. 13. 6. Incidental Hosts:

In the scrub-typhus cycle, humans become incidental hosts. Humans can become infected with *Orientiatsutsugamushi* when infected chiggers bite and consume human blood through their saliva.

1. 13. 7. Clinical Manifestation:

Human scrub typhus is the infection's clinical manifestation, characterized by fever, headache, muscle soreness, and eschar formation at the bite site.

2. Epidemiology of scrub typhus:

Highly endemic to the "tsutsugamushi's triangle," scrub typhus inhabits an area of over 8 million square kilometres that stretches from far eastern Russia in the north to Pakistan in the west, Australia in the south, and Japan in the north. Several studies have reported that the tsutsugamushi's triangle has been the source of most scrub typhus cases. However, enteric studies understand that scrub typhus is spreading over the world and is no longer restricted to the Tsutsugamushi's triangle. Scrub typhus has been found to be primarily acquired by a number of factors, including the presence of woodpiles, scrub vegetation, and cattle near residential areas(30). According to a study, environmental factors appear to be the primary cause of scrub typhus occurrence, rather than behavioural or demographic factors. Scrub typhus is more common in areas where there is scrub vegetation near homes, cooking outdoors, domesticating pets, and living close to a body of water. The two ways that Orientals tsutsugamushi spreads are through the eggs that a female transmits to her offspring and through the passage of mite larvae, nymphs, and adults. Both techniques are classified as vertical transmission, and there is currently no proof of horizontal transmission of *O. tsutsugamushi*—where a mite obtains Oriental from an infected host and then spreads the infection to other hosts—according to available data. The majority of the cases in the Asia-Pacific area were in China's eastern, southeast, and southwest coastal regions. The age group of 60–69 years old in China had cumulative incidences, with a maximum observed in June and July. Without any variation in incidence between genders, the majority of cases (23.36%) were reported in the 50–60-year-old age group. Unlike China, Japan experienced its greatest scrub typhus outbreaks in November. The distribution of ages differed from China as well, with the majority of cases (62%) occurring in the 51–75 age range. In South Korea, scrub typhus is one of the most common rickettsial illnesses, accounting for 27.7–51% of all cases of acute febrile illness. Notably, there is a gender disparity in the disease, with more female cases than males (Chang 1995). Taiwan has reported a higher female-to-male ratio, with a higher incidence in the 50–60-year-old age group. Japan's gender equality ratio is comparable to that of Taiwan. In line with Japan, October and November saw the highest frequency of disease occurrence among patients aged 60–69 in South Korea (27.48%). **Scrub typhus is a disease that still goes undiagnosed in India, affecting people from the south to the northeast and northwest.** August through November was noted as the peak season for disease occurrence, with rural areas of India being the primary affected(31). According to sera-epidemiological research, scrub typhus is more common in Thailand, where cases significantly increased between the 1980s and 2000s. In contrast to China and Japan, where disease incidence is higher (22.3%) among patients aged 30-39, Thailand has a reported male-to-female gender ratio of 2:1. The disease was first discovered during the Vietnam War in Vietnam and other Southeast Asian nations, with army personnel accounting for the majority of cases diagnosed in the 1960s. However, the illness was disregarded until the early 21st century, which resulted in gaps in the literature and inadequate data gathering for epidemiological research (Deller and Russell 1967; Thiebaut et al. 1997). Still, new research from Vietnam indicates that the cumulative incidence of scrub typhus was 1.1% in the general population and 3.5% in patients who were hospitalized, with no discernible difference between rural and urban areas. Scrub typhus is also endemic in a small number of other countries that are part of the "tsutsugamushi's triangle," which also includes Indonesia, Malaysia, the Philippines, Pakistan, Far East Russia, and Australia. These countries were first identified as "coastal fever" in 1913 and subsequently as scrub typhus. Scrub typhus was found to be present outside of the traditional endemic areas in both serological and molecular studies. This suggests that the disease is now a global problem and is no longer limited to the "tsutsugamushi's triangle" as it was previously thought. In order to control the disease, epidemiological studies are crucial for comprehending how the disease is distributed across various age, gender, and geographic

groups. The emergence of scrub typhus outside of its native range raises the prospect of undiscovered Oriental species and new scrub typhus vectors(32).

2.1. Endemic Regions and Affected Populations:

The bite of an infected chigger (a larval mite) can expose humans to the bacterial disease scrub typhus, which is caused by *Orientia tsutsugamushi*. The term "endemic regions" refers to specific areas around the Asia-Pacific rim where the disease is predominantly prevalent. These domains consist of:

2. 1. 1. Southeast Asia:

Scrub typhus is known to have endemic areas in nations like Vietnam, Laos, Thailand, Malaysia, Indonesia, and Cambodia.

2. 1. 2. South Asia:

Scrub typhus also affects areas of Bhutan, India, Nepal, and parts of Sri Lanka.

2. 1. 3. East Asia:

The scrub typhus cases have been reported in South Korea, Japan, and parts of China.

2. 1. 4. Pacific Islands:

These are reported cases of scrub typhus on several Pacific islands, including Papua New Guinea and parts of northern Australia.

People who live in rural or agricultural areas with close proximity to vegetation and environments that support chigger habitats are usually the most affected populations. Workers in agriculture, military personnel, and people who go outside in these endemic areas are more likely to get scrub typhus. Poor sanitation and rural settings are frequently linked to the disease(33).

Scrub typhus is characterized by a dark, scab-like lesion at the site of the chigger bite, fever, headache, and muscle soreness. Scrub typhus can cause more serious consequences, like organ failure, if it is not treated.

Wearing protective clothes, applying insect repellents, and staying away from places with dense vegetation and potential chigger habitats are some preventive measures. Scrub typhus must be managed, and serious consequences can be avoided with early diagnosis and appropriate antibiotic treatment.

2. 2. Seasonal Patterns and Environmental factors influencing transmission:

2. 2. 1. Seasonal Patterns:

Scrub typhus often exhibits a seasonal occurrence, with an increased number of cases during specific times of the year.

The disease is commonly associated with the rainy season or monsoon period in endemic regions. This is because chiggers—mites that spread the bacteria that cause scrub typhus—need this period of increased vegetation and humidity to breed and survive(34).

2. 2. 2. Environmental Factors:

2. 2. 2. 1. Vegetation and Habitat:

Dense vegetation, tall grasses, and shrubs are associated with scrub typhus. In these environments, chiggers live as larvae and attach themselves to people when they are outdoors.

2. 2. 2. 2. Rural and Agricultural Settings:

In agricultural and rural locations where people have close contact with plants, the disease is more common. The risk of exposure is increased by farming operations and living close to fields.

2. 2. 2. 3. Altitude:

Altitude has the potential to affect scrub typhus incidence. Certain elevations in some regions may have a higher prevalence of the disease.

2. 2. 2. 4. Human Behaviour:

2. 2. 2. 5. Outdoor Activities:

Participating in outdoor pursuits like farming, camping, or trekking within endemic areas raises the probability of encountering chigger-infested environments.

2. 2. 2. 6. Clothing Practices:

Wearing protective apparel, like long sleeves and pants, can lower the chance of chigger bites and, in turn, the spread of scrub typhus.

2. 2. 2. 7. Climate Conditions:

2. 2. 2. 8. Humidity:

High humidity is ideal for chiggers. Chiggers are more likely to survive and be active during the rainy season because it produces a humid atmosphere.

2. 2. 2. 9. Temperature:

Scrub typhus prevalence in endemic areas can be influenced by temperature range; warmer temperatures are frequently more favourable to chigger activity.

2.3. Clinical presentation:

There is a wide range of clinical manifestations of scrub typhus that can differ in intensity. The bacterium *Orientia tsutsugamushi*, which causes the disease, is contracted by humans through the bite of an infected chigger.

2. 3. 1. Incubation Period:

Scrub typhus usually takes six to twenty-one days to incubate after a chigger bite.

2. 3. 2. Onset of Symptoms:

Scrub typhus is characterized by a sudden onset of fever, which is often the first symptom of the illness.

2. 3. 3. Fever:

A fever that is elevated is a typical and noticeable symptom. Chills may accompany the fever, which may last for several days.

2. 3. 4. Headache:

Another common symptom that people with scrub typhus report is headaches.

2. 3. 5. Muscle and Joint Pain:

Joint pain and muscle aches (malign) are common in patients, which add to their overall discomfort.

2. 3. 6. Eschar Formation:

An eschar, or dark, scab-like lesion, forms at the site of the chigger bite and is a distinctive feature of scrub typhus. Usually painless, the eschar may go unnoticed.

2. 3. 7. Lymphadenopathy:

Lymphadenopathy—swollen, tender lymph nodes—usually appears in the vicinity of the chigger bite site.

2. 3. 8. Rash:

A rash might appear, but it doesn't always. When it does appear, the rash is typically maculopapular, or small, raised red spots, and it can spread to other body areas.

2. 3. 9. Gastrointestinal Symptoms:

It is possible to experience gastrointestinal symptoms like nausea, vomiting, diarrhoea, and abdominal pain.

2. 3. 10. Respiratory Symptoms:

Respiratory symptoms like coughing and breathing difficulties may appear in severe cases.

2. 3. 11. Central Nervous System Involvement:

Some people may develop neurological symptoms such as drowsiness, confusion, and, in extreme situations, meningitis.

2. 3. 12. Complications:

Scrub typhus can cause serious complications, including organ failure, if it is not treated promptly and appropriately.

3. Diagnosis:

3. 1. Clinical Evaluation:

Numerous non-specific symptoms, such as fever, headaches, muscle aches, and a distinctive eschar—a dark, scab-like lesion—at the site of the chigger bite, are indicative of scrub typhus.

The clinical presentation can differ, and symptoms can sometimes mimic those of other feverish illnesses. A thorough medical history is therefore essential, including details about any recent travel or exposure to endemic areas(35).

4. Laboratory Tests:

4. 1. 1. Serological Tests:

To diagnose scrub typhus, pathological testing is frequently employed. Common techniques to identify antibodies (IgM and IgG) against the causative bacterium, *Orientia tsutsugamushi*, are the enzyme-linked immunosorbent assay (ELISA) and the indirect immunofluorescence assay (IFA).

4. 1 . 2. Polymerase Chain Reaction (PCR):

Blood sample bacteria can be identified genetically using PCR tests. With this approach, the infection can be confirmed more precisely.

4. 1. 3. Weil-Felix Test:

Despite being less precise, this test could be applicable in certain contexts. Finding agglutinating antibodies against specific *Proteus* species that cross-react with *Orientia tsutsugamushi* is the method involved.

4. 1. 4. Imaging Studies:

Imaging tests, like chest X-rays, might be performed in extreme situations to determine the degree of organ involvement.

4. 1. 5. Clinical Criteria:

Clinical criteria may also be used to make a diagnosis, particularly in situations with limited resources where laboratory testing may not be easily accessible. Symptoms, exposure history, and treatment response are all taken into account.

The timely and efficient treatment of scrub typhus depends on an early diagnosis. Serious side effects, such as organ failure, may result from the illness if treatment is stopped. Because of this, medical professionals frequently start treating patients based on clinical suspicion even before they have access to confirmatory test results (36).

4. 1.6. Treatment and management:

Treatment:

4. 1. 6. 1. Antibiotics:

Giving antibiotics, usually doxycycline is the main and most successful treatment for scrub typhus. In instances where doxycycline is contraindicated or poorly tolerated, alternative antibiotics like azithromycin or chloramphenicol might be employed.

4. 1. 6. 2. Timing is crucial:

A good result depends on starting antibiotic therapy as soon as possible. Treatment setbacks may result in more serious side effects.

5. Management:

5. 1. 1. Supportive Care:

To control symptoms and avoid complications, patients with scrub typhus may need supportive care. Pain relief, maintaining a fever, and getting enough water are examples of supportive measures.

5. 1. 2. Monitoring:

Vital signs such as blood pressure, heart rate, and temperature should be regularly monitored. Evaluation of the infection's severity may involve laboratory testing, such as liver function tests and complete blood counts.

5. 1. 3. Complication Management:

Acute respiratory distress syndrome (ARDS), Meningoencephalitis, and organ failure are among the consequences of severe scrub typhus. Interventions specific to the management of complications may include support for intensive care.

5. 1. 4. Preventive Measures:

Avoiding chigger exposure is essential to preventing scrub typhus. This entails donning protective gear, applying insect repellents, and staying away from places with a lot of vegetation where chiggers are frequently encountered.

5. 1. 5. Public Health Measures:

Initiatives aimed at improving public health may involve the use of health education campaigns to increase knowledge about preventive measures and vector control programs to lower the prevalence of mites.

5. 2. Prevention and Control

5. 2. 1. Personal Protection

5. 2. 2. Wearing Protective Clothing

Wearing closed shoes, socks, long sleeves, and long pants when venturing into densely vegetated areas can help ward off chigger bites.

5. 2. 3. Insect Repellents:

Applying insect repellents containing DEET (N, N-diethyl-meta-polyamide) to exposed skin can deter chiggers.

5. 2. 4. Permethrin-Treated Clothing

Additional protection can be obtained by treating clothing with the insect repellent permethrin.

5.3. Avoiding High-Risk Areas:

5.3.1. Vegetation Awareness:

Use caution when in areas that have bushes, tall grass, and other vegetation that is known to harbour chiggers.

5.3.2. Outdoor Activities:

Reducing outdoor activity in high risk- areas when mite activity is at its highest.

5.3.3. Environmental Measures:

5.3.3.1. Vector Control:

Using acaroids, or chemicals that kill mites, in impacted areas as one method of managing mite populations.

5.3.3.2. Vegetation Management:

Removing vegetation and keeping the surroundings tidy to lessen chigger-friendly habitats.

5.4. Public Health Education:

5.4.1. Awareness Campaigns:

Establishing community education programs about the dangers of scrub typhus, the value of personal protection, and the early detection of symptoms.

5.4.2. Healthcare Provider Training:

Ensuring that medical professionals are knowledgeable about scrub typhus in order to diagnose and treat patients on time.

6. Prompt Diagnosis and Treatment:

6.1.1. Healthcare Access:

Ensuring prompt medical attention is sought by those exhibiting symptoms suggestive of scrub typhus.

6.1.2. Timely Antibiotic Treatment:

To prevent complications, give the right antibiotics as soon as a diagnosis is made.

7. Surveillance and Reporting:

7.1.1. Disease Monitoring:

Establishing surveillance systems to track scrub typhus distribution and prevalence.

7.1.2. Reporting Cases:

Ensuring that healthcare facilities promptly report confirmed cases to public health authorities for timely response.

8. Research and Development:

8.1. Vaccine Development:

Supporting studies aimed at creating a scrub typhus vaccine that works.

8.1.1. Vaccine Development and Challenges:

8.1.2. Target Antigen Identification:

The main goal of research is usually to find antigens, or proteins, on the surface of the bacterium *Orientia tsutsugamushi* that can trigger an immune response. There's a chance that a vaccine will contain these antigens.

8.1.3. Preclinical Testing:

Preclinical testing is done on identified potential antigens in lab and animal models to evaluate their safety, immunogenicity, and effectiveness.

8.1.4. Clinical Trials:

Promising vaccines proceed to human clinical trials if positive preclinical study results are obtained. To assess safety, dosage, and effectiveness in broader populations, clinical trials are conducted in three stages.

8.1.5. Regulatory Approval:

Obtaining regulatory approval requires completing clinical trials successfully and satisfying safety and efficacy standards. The vaccine is evaluated by regulatory agencies such as the European Medicines Agency (EMA) or the U.S. Food and Drug Administration (FDA) prior to its commercialization.

9. Challenges in Scrub Typhus Vaccine Development:

9.1.1. Complexity of the Bacterium:

Orientia tsutsugamushi is a multistrain complex bacterium. It can be difficult to develop a vaccine that offers broad protection against various strains.

9.1.2. Limited Understanding of Immune Response:

Uncertainty exists regarding the specific immune response needed to ward off scrub typhus. It is critical to determine the most potent antigens and develop a vaccine that elicits a robust and long-lasting immune response.

9.1.3. Lack of Standardized Animal Models:

It is difficult to find appropriate animal models that faithfully simulate human infection. It can be challenging to forecast the effectiveness of vaccines because certain animals exhibit distinct disease manifestations compared to humans.

9. 1. 4. Geographical Variation:

Different strains are more common in different regions. Due to regional differences in strain prevalence, a vaccine that is effective in one area might not offer the same level of protection in another.

9. 1. 5. Limited Commercial Interest:

Since scrub typhus is mostly endemic in particular areas, pharmaceutical companies may not be very interested in investing in the development of a vaccine. This may have an effect on research funding and resources.

9. 1. 6. Ethical and Logistical Challenges in Clinical Trials:

Clinical trial conduct for infectious diseases poses ethical problems, and in areas where scrub typhus is endemic, logistical problems could surface. Clinical trial success depends on adhering to the right ethical standards and overcoming practical obstacle(37).

10. Conclusion:

In summary, combating scrub typhus necessitates a multimodal strategy that incorporates ongoing research projects, clinical care, and public health initiatives. For effective treatment with antibiotics like doxycycline, a timely diagnosis made possible by improved diagnostic tools is essential. In endemic areas, prevention strategies include community education, vector control, and personal protective measures to lower the risk of chigger exposure. Despite the advancements, issues like antibiotic resistance and the lack of a generally accepted vaccine still exist. Effective disease management requires cooperation between communities, researchers, and healthcare professionals. Prevention of this neglected tropical disease is largely dependent on ongoing research, particularly on vaccine development. We can work toward a future where scrub typhus is better controlled and its impact on impacted populations is greatly lessened by tackling these issues and putting comprehensive strategies into place.

Acknowledgements: All authors expressed the heartfelt thanks to the management of Raghavendra Institute of Pharmaceutical Education and Research (RIPER) – Autonomous for providing data bases, collection of information on recent developments of scrub typhus.

11. References:

1. Chakraborty S, Sarma N. Scrub typhus: an emerging threat. *Indian Journal of Dermatology*. 2017;62(5):478.
2. Mahajan S. Scrub typhus. *JAPI*. 2005;53(955):269.
3. Traub R, Wisseman Jr CL. Ecological considerations in scrub typhus: 1. Emerging concepts. *Bulletin of the World Health Organization*. 1968;39(2):209.
4. Luce-Fedrow A, Lehman ML, Kelly DJ, Mullins K, Maina AN, Stewart RL, et al. A review of scrub typhus (*Orientia tsutsugamushi* and related organisms): then, now, and tomorrow. *Tropical medicine and infectious disease*. 2018;3(1):8.
5. Jagadeesan S, Patel P, Kushwaha P, Patidar N. Scrub typhus fulminating as liver failure: A rare report. *Journal of Vector Borne Diseases*. 2023;60(3):333-5.
6. Cho H-W, Chu C. The geographical and economical impact of scrub typhus, the fastest-growing vector-borne disease in Korea. *Osong Public Health and Research Perspectives*. 2013;4(1):1.

7. Organization WH, Research SPf, Diseases TiT, Diseases WHoDoCoNT, Epidemic WHO, Alert P. Dengue: guidelines for diagnosis, treatment, prevention and control: World Health Organization; 2009.
8. Janardhanan J, Trowbridge P, Varghese GM. Diagnosis of scrub typhus. Expert review of anti-infective therapy. 2014;12(12):1533-40.
9. Richards AL, Jiang J. Scrub typhus: historic perspective and current status of the worldwide presence of *Orientia* species. Tropical medicine and infectious disease. 2020;5(2):49.
10. Dorji K, Phuentshok Y, Zangpo T, Dorjee S, Dorjee C, Jolly P, et al. Clinical and epidemiological patterns of scrub typhus, an emerging disease in Bhutan. Tropical Medicine and Infectious Disease. 2019;4(2):56.
11. Singer M. Introduction to syndemics: A critical systems approach to public and community health: John Wiley & Sons; 2009.
12. Higuchi H. Natural history of Japanese birds. Heibonsha, Tokyo. 2014.
13. Park S-W, Ha N-Y, Ryu B, Bang JH, Song H, Kim Y, et al. Urbanization of scrub typhus disease in South Korea. PLoS neglected tropical diseases. 2015;9(5):e0003814.
14. Park HS, Lee JH, Jeong EJ, Kim JE, Hong SJ, Park TK, et al. Rapid and simple identification of *Orientia tsutsugamushi* from other group rickettsiae by duplex PCR assay using *groEL* gene. Microbiology and immunology. 2005;49(6):545-9.
15. Kweon S-S, Choi J-S, Lim H-S, Kim J-R, Kim K-Y, Ryu S-Y, et al. A community-based case-control study of behavioral factors associated with scrub typhus during the autumn epidemic season in South Korea. The American journal of tropical medicine and hygiene. 2009;80(3):442-6.
16. Iqbal MZ. Serodiagnosis of Scrub Typhus in a Tertiary Care Hospital: Rajiv Gandhi University of Health Sciences (India); 2018.
17. Brig Amar Cheema V. The Crimson Chinar: The Kashmir Conflict: A Politico Military Perspective: Lancer Publishers; 2015.
18. Varghese GM, Janardhanan J, Trowbridge P, Peter JV, Prakash JA, Sathyendra S, et al. Scrub typhus in South India: clinical and laboratory manifestations, genetic variability, and outcome. International Journal of Infectious Diseases. 2013;17(11):e981-e7.
19. Raghavendra K, Velamuri PS, Verma V, Elamathi N, Barik TK, Bhatt RM, Dash AP. Temporo-spatial distribution of insecticide-resistance in Indian malaria vectors in the last quarter-century: need for regular resistance monitoring and management. Journal of vector borne diseases. 2017;54(2):111.
20. Poonia T, Singh N, Garg M. Contamination of Arsenic, Chromium and Fluoride in the Indian groundwater: a review, meta-analysis and cancer risk assessment. International Journal of Environmental Science and Technology. 2021;18:2891-902.
21. Wangrangsimakul T. Scrub typhus in northern Thailand: University of Oxford; 2021.
22. Kelly DJ, Fuerst PA, Ching W-M, Richards AL. Scrub typhus: the geographic distribution of phenotypic and genotypic variants of *Orientia tsutsugamushi*. Clinical infectious diseases. 2009;48(Supplement_3):S203-S30.
23. Tantibhedhyangkul W, Amara AB, Textoris J, Gorvel L, Ghigo E, Capo C, Mege J-L. *Orientia tsutsugamushi*, the causative agent of scrub typhus, induces an inflammatory program in human macrophages. Microbial pathogenesis. 2013;55:55-63.
24. Zaman K. Scrub typhus, a salient threat: Needs attention. PLOS Neglected Tropical Diseases. 2023;17(6):e0011427.
25. Shil R. Scrub Typhus—An Emerging Public Health Threat in Northern India. Saudi J Med Pharm Sci. 2023;9(10):708-12.
26. Ali A, Waris A, Khan MA, Asim M, Khan AU, Khan S, Zeb J. Recent advancement, immune responses, and mechanism of action of various vaccines against intracellular bacterial infections. Life Sciences. 2022:121332.
27. Traub R, Wisseman Jr CL. The ecology of chigger-borne rickettsiosis (scrub typhus). Journal of medical entomology. 1974;11(3):237-303.
28. Elliott I, Pearson I, Dahal P, Thomas NV, Roberts T, Newton PN. Scrub typhus ecology: a systematic review of *Orientia* in vectors and hosts. Parasites & vectors. 2019;12(1):1-36.
29. Gakpey EE. Understanding the Dynamics of Climate Change, Migration and Conflict: The Case of Bangladesh. 2022.
30. Govindarajan R, Rajamannar V, Kumar A, Samuel PP. Current status of mites and mite-borne diseases in India. Journal of Vector Borne Diseases. 2023;60(1):1-10.
31. Gupta S. Diversity, biology and pathogenesis of *Orientia tsutsugamushi* (Scrub typhus pathogen): A general.
32. Lamichhane P, Pokhrel KM, Alghalyini B, Zaidi ARZ, Alshehry MZ, Khanal K, et al. Epidemiology, clinical characteristics, diagnosis, and complications of scrub typhus infection in Nepal: a systematic review. Annals of Medicine and Surgery. 2023;85(10):5022-30.

33. Machtinger ET, Weeks EN, Geden CJ, Lacher E. Pests that develop and live on horses (ectoparasites). *Pests and parasites of horses: Wageningen Academic*; 2022. p. 141-97.
34. McCown ME. *Zoonotic and Infectious Vector/Tick Borne Pathogen Surveillance in Military Working Dogs, Police Working Dogs and Canines in Colombia, South America: University of Florida*; 2022.
35. Nair LVK, Timmalsugur RK, Mangal V. Scrub Typhus Meningoencephalitis: An Uncommon Cause of Acute Febrile Encephalopathy in the Elderly. *Journal of Marine Medical Society*. 2022;24(Suppl 1):S101-S3.
36. Geisler WM, Uniyal A, Lee JY, Lensing SY, Johnson S, Perry RC, et al. Azithromycin versus doxycycline for urogenital *Chlamydia trachomatis* infection. *New England Journal of Medicine*. 2015;373(26):2512-21.
37. Patil SD, Raut MK, Rathod NP. Quality Control And Quality Assurance In Pharmaceuticals. *Quality Control And Quality Assurance In Pharmaceuticals*. 2023;1(10):1-.

UNDER PEER REVIEW