

Molecular diagnosis with histological study of *Eimeria* spp. (*Camelus dromedaries*) at AL-Najaf AL-Ashraf provinces

Abstract

Many nations rely heavily on the export of Old World camels for their economic needs. They are employed in camel transportation, the production of food such as meat and milk, and racing

Eimeria affects a wide variety of domestic including camels and wild animals,. The aims of this study was to distribution and histological effected of *Eimeria* spp. camels (*Camelus dromedaries*) using Molecular diagnosis with histological study. 100 fecal samples of camels (*Camelus dromedaries*) were examined by conventional PCR from different reign at AL-Najaf AL-Ashraf province of Iraq, from Novmber 2023 to end of April 2024. A total infection rate with *Eimeria* spp was 57% (57/100) . male camels recorded higher than female 63.2% (24/38) . The age group of male at (<5 years) show the higher rate of infection 76.7% (23/30) while the female recorded 80.8% (9/11) in age (10 years \geq) as a higher infection rate. district Al- Manathera was recorded 71.4% (12/29) with significant differences at

($P \leq 0.05$) compare with anther district of AL-Najaf AL-Ashraf provinces.

A total of 20 samples collected from small intestine of slaughtered camels were collected. Histopathological analysis of the infected tissues revealed granulomatous, hemorrhagic, and chronic enteritis; in some cases, the intestinal glands have a rose-shaped appearance, while in other cases, there is a severe infiltration of chronic inflammatory cells with no intestinal gland structure; hypereosinophilia and various developmental stages of *Eimeria* are also present.

Keywords: Food, *Eimeria*, infection , *Camelus dromedaries* molecular diagnosis , Histopathologica .

Introduction

The camel is a significant and unusual mammal that has adapted to survive and procreate in the intense heat and dryness of the desert. a significant food source, transit for African and Asian desert nomads. One of the best kinds of milk is camel's milk. because it has more iron and vitamin C than milk from cows. *Eimeria* spp is one of Parasitic infections, particularly gastrointestinal which have had impact on camels development and output. The economic harm they cause, including the mortality of animals and the diminution of livestock products, is significant, particularly in young animals (Kamal *et al.*, 2024). The feces of both symptomatic and carrier animals contain oocysts shed by host-specific protozoan parasites called coccidia. These carriers can transmit infection through water and food sources

(Jilo *et al.*, 2022). Coccidiosis is an obligatory illness that causes hypertrophy and death in parasitized cells (Hassan *et al.*, 2024) Coccidia can cause injury to host's intestinal cells cause anemia and a loss of Electrolytes (Murshed *et al.*, 2024)

When animals consume contaminated feed or water, the parasite injures intestinal cells, causing diarrhea and hematochezia (Utebaeva *et al.*, 2021). The great resilience of oocysts to environmental conditions is a crucial aspect of the infection's epidemiology (Gao *et al.*, 2024).

Eimeria's classification was based on the morphological features of both sporulated and non-sporulated oocysts (Hegazi *et al.*, 2023).

Materials and Methods

A total of (2) grams fecal samples were taken from (100) camels (38 male, 62 female) of various ages from certain region in Al-Najaf, Al-Ashraf, between November 2023 and the end of April 2024. Fecal samples obtained in sterile plastic containers were firmly sealed, and safety precautions such as disposable gloves were removed owing to consecutive numbers. All sample information, including age, gender, and date, were sent in a cooling box to a laboratory of Parasitology / College of Veterinary Medicine- University Al-Qasim Green University / Iraq for Molecular diagnosis.

20 samples were collected from intestine of slaughtered one humped camels at Al-Najaf, Al-Ashraf. Tissue samples were performed within 1 to 5 hours after death. Samples were taken from intestines which showed gross lesion and used for histopathological examination.

Genomic DNA Extraction

Genomic DNA from feces samples were extracted by using Presto stool DNA Extraction Kit, (Geneaid Biotech /China) and done according to company instructions.

Polymerase chain reaction (PCR)

The PCR technique was performed for detection of *Eimeria spp.* in feces samples in camels based on 18S small subunit ribosomal RNA gene, This method was carried out according to method described by (Zainabet *al.*, 2016)

Preparation of histological sections:

Samples from various regions of the small intestine (duodenum, colon, and jejunum) were collected and fixed in 10% neutral buffered formalin solution, then washed, dehydrated in various grades of ethyl alcohol, clarified in methyl benzoate, and embedded in paraffin wax. Blocks were handled according to conventional processes .Used hematoxylin-eosin to stain 5- μ m thick sections for microscopy analysis according to (Spenceret *al.*, 2012).

Statistical analysis

The Chi-square test was used to detect the association between parasite infection rate and each animal (camel). P-values < 0.05 were regarded statistically significant using SPSS (statistical package for social sciences) version 26 (SPSS, 2020). User Guide Statistic Version, 26th Edition. SPSS, the statistical tool for social science, user handbook statistical version, sixth edition.

Results

The total rate of infectionwith *Eimeria spp.* in camels (*Camelus dromedaries*)by Molecular diagnosisfrom different districtat AL-Najaf AL-Ashraf province of Iraq, during Novmber 2023to end of April 2024. A total infection rate with *Eimeriaspp* was 57% (57/100) (Table 1).

Table (1):Rate of infection with *Eimeria spp.* In camels by using Molecular diagnosis.

Host	Fecal sample examined	Molecular diagnosis	
		NO. of positive	Percentage%
Camel	100	57	57

Infection

rate with *Eimeria spp.* in camels according to sex byMolecular diagnosis:-

According to sex male and female recorded 63.2% (24/38) and 53.2% (33/62) rate of infection respectively. with no significant differences ($p \leq 0.05$) (Table 2).

Table (2):Infection rate with *Eimeria spp.* in camels according to sex.

Sex	No. of samples examined	No. of positive	Percentage (%)

Male	38	24	63.2
Female	62	33	53.2
Total	100	57	57
X ²	0.948229		
P value	0.330171		

Nosignificant differences at (P≤0.05).

Rate of infection with *Eimeria* spp. in camels male according to age groups

Infection with *Eimeria* spp. was recorded Highest rate in camles 76.7% (23/30), whilerecorded in groups (10 years ≥) 0% (0/2) with significant differences (p≤0.05) (Table 3).

Table (3): Rate of infection with *Eimeria* spp. in camels male according to age

exam.samples		No.	%oftotal
5years≤	30	23	76.7
5-10years	6	1	16.7
10years≥	2	0	0
Total	38	24	63.2
X ²	11.354762		
Pvalue	0.003423*		

*: significant differences at (P≤0.05).

Rate of infection with *Eimeria* spp. in camels fmale according to age groups

Highest rate of infection with *Eimeria* spp. was recorded in female 81.8 % (9/11), while lowest recorded in groups (5 years≤) 10 % (2/20) with significant differences (p≤0.05)(Table 4).

Table (4):Rate of infection with *Eimeria* spp. in camels fmale according to age

Age(year)	No.ofthe exam.samples	Positivesamples	
		No.	%oftotal
5years≤	20	2	10
5-10years	31	22	71
10years≥	11	9	81.8
Total	62	33	53.2
χ^2	22.541921		
Pvalue	0.000013**		

** : significant differences at ($P \leq 0.05$).

Rate of infection with *Eimeria* spp. in camels according to the district in AL-Najaf AL-Ashraf

The rate of infection with *Eimeria* spp in camel samples at three district as following AL-Manathera 71.4% (25/35), Slaughter house 55.6% (20/36,) and AL-Meshkab 41.4% (12/29) respectively with significant differences ($p \leq 0.05$) (Table 5) .


Table (5): Rate of infection with *Eimeria* spp. in camels according to the district in AL-Najaf AL-Ashraf

District	No.ofthe exam. samples	Positivesamples	
		No.	%oftotal
Slaughterhouse	36	20	55.6
AL-Manathera	35	25	71.4
AL-Meshkab	29	12	41.4
Total	100	57	57
χ^2	5.890539		
Pvalue	0.05*		

*: significant differences at ($P \leq 0.05$).

Conventional PCR product analysis:

Genomic DNA was obtained from camel fecal samples and subjected to molecular analysis by using PCR for detection of small subunit ribosomal RNA gene. Specific primers were used to identify the species of *Eimeria*. PCR of 100 samples was employed in the study exhibited a distinct band of (456 bp) from the PCR product on the agarose gel.



Figure(1) PCR products of amplification of 18S rRNA gene for detection of *Eimeria* species. Size of PCR product: 456 bp. The gel was 1.5% and the DNA dye is RedSafe (Intron, Korea). Electrophoresis conditions: V: 95, Time: 45 minutes. M: DNA ladder

DNA sequencing results

7 out of 57 samples from were positive by using Conventional PCR and sequencing depots in NCBI Genbank data base and DDBJ to get accession number codes (PQ100675, PQ111747, PQ129311,PQ129315, PQ129400. PQ120694 and PQ113350) for of local *Eimeria* species at the first time in Iraq.

Phylogenetic confirmative detection

Specific phylogenetic confirmative detection of local *Eimeria* species was performed by using the phylogenetic tree analysis and compared with NCBI-BLAST *Eimeria* species (figure2).




Figure (2): Specific phylogenetic confirmative detection of local *Eimeria* species

Histological change results

They demonstrated villous atrophy, degeneration, necrosis, and denudation. The mucosal glands in the afflicted gut were likewise found to be vacuolated, with epithelial cell disintegration. Occasionally, the crypt epithelium detached from the basement membrane. Catarrhal colitis, goblet cell hyperplasia, and isolated chronic inflammation in the mucosa with large lymphocytic infiltration were clearly visible, as well as chronic enteritis, hemorrhagic enteritis, and granulomatous enteritis with goblet cell proliferation of the intestinal gland. Furthermore, the infiltration of chronic inflammatory cells between intestinal glands was found. Instead of the large intestine, the lamina propria of the small intestine had cccidian stages. Many nuclei grouped together in blastophores. The parasites are found intracellularly in sizable parasitophorous vacuoles; they are also found macrogamontly in the lamina propria, where they have completely destroyed the intestinal glands and infiltrated the cells responsible for chronic inflammation. microgamont with several nuclei (figure3).




Figure (3): Histopathological section of camel intestine infected with *Eimeria* spp. showing different development stage of *Eimeria* (red circle) with inflammatory cells infiltration (red arrow), also atrophy and sloughing of intestinal villi (black arrow) necrosis in intestinal gland (black circle), with edema among the muscularis layer (black arrowhead) (H & E stains, 200x).




Figure (4): High magnification for inflammatory cells infiltration showing presence of eosinophile (red arrow), macrophage (black arrow) and lymphocyte (yellow arrow) H & E stains, 400x.




Figure (5): Histopathological section of camel intestine infected with *Eimeria* spp showing presence of mature macrogamonts inside the intestinal glands (yellow arrow) (H &E stains, 400x).

Discussion

The large class of parasites known as coccidia has to be present within the cell. Numerous cattle species are impacted by the economically significant sickness known as coccidiosis.

(Ashfaq *et al.*, 2023). The symptoms of coccidiosis vary from indifference and mild, transient diarrhea to severe instances that include large amounts of black, bloody diarrhea and, in the worst situations, fatality. In order to handle coccidiosis in camels in an economical and efficient manner, a thorough understanding of the *Eimeria* species involved is necessary. (El-Bahyet *et al.*, 2023).

Thus, identifying the frequency and variety of *Eimeria* species was the goal of the current study. Prior to this investigation, *Eimeria* species were thought to be harmful to newborn camel calves.

(Hussein *et al.* 1987). The age and sex of the camels in the current investigation on camel coccidiosis did not significantly affect the prevalence. These results disagreed with those of Yakhchali and Cheraghi (2007). Due to their greater sensitivity to infection, the breeding and overcrowding systems seen in their various qualities, and their immature immune system development as compared to older ruminants, young ruminants are more prone to illness than older ones. (Safaa and May ,2020).

Concurring to a few thinks about the variability in prevalence and distribution of coccidiosis may be due to variations in management and hygienic conditions, temperature, agro ecology,

environment, weather conditions, host immune status, sample size, sampling period and breed susceptibility to coccidia in different areas

The ileum, colon, and jejunum were the most frequently afflicted tissues in this research with chronic enteritis; the discovery of gigantic schizonts in the lamina propria with total destruction of intestinal glands and infiltration of chronic inflammatory cells was consistent with (Kheirandish *et al.*, 2012). Large parasitophorous vacuoles containing the parasites are seen inside cells. Finding the macrogamont within the lamina propria and total annihilation of gut glands containing cells involved in chronic inflammation invasion, multi-nucleated microgamont; There are debris strands in the parasitophorous vacuole. was comparable to the outcomes of (Dubey *et al.*, 2018).

The intestinal mucosa included *E. cameli* oocytes, meronts, and macrochizonts, which are various developmental stages. These phases belonged to *Eimeria* spp, as Immature oocysts of *Eimeria* spp were discovered within the Intestinal epithelial mucosa. To our knowledge, the formation of *Eimeria* spp oocyte wall is unclear. In *Eimeria* species, the oocyst wall is initiated by the secretions of WFB. There are two types of WFB in most *Eimeria* species, type 1 and type 2 (Ferguson *et al.*, 2003)

As the macrogamont matures, the WFB develops the oocyst wall (Ferguson *et al.*, 2003). Furthermore, there are other veil-forming (VFB); in certain coccidian parasites, they generate a veil-like outer layer on the oocyst wall; once the Oocysts are expelled in feces, and the wall has vanished.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

1.

References

- Al-Kaabi, N.A. (2009) Epidemiological and diagnostic study of coccidiosis in sheep of Diwaniya. M.Sc. thesis. University of Al-Qadisiya, College of veterinary Medicine. Pp.; 33-38.
- Ashfaq, K., Asghar, A. Y., Hashmi, S. S., & Abbas, A. (2023). Bovine Coccidiosis A formidable challenge to cattle industry. *International Journal of Research and Advances in Agricultural Sciences*, 2(2), 34-42.
- Coles, E. H. (1986) Vet. Clinical Pathology 4th ed., W. B. Saunders Company, Philadelphia. Pp.:375-379
- Dubey, J.P., Rolf, K., Schuster, and Joerg K.(2018)Gametogony of *Eimeria cameli* in the small intestine of one-humped camel (*Camelus dromedaries*, *Parasitology Research* 117:3633–3638
- El-Bahy, M. M., Kamel, N. O., Auda, H. M., and Ramadan, R. M. (2023). A smart economic way to control camel parasites and improve camel production in Egypt. *Experimental Parasitology*, 255, 108650.
- Ferguson, D.J.P., Belli, S.I., Smith, N.C., and Wallach,M.G.(2003). The development of the macrogamete and oocyst wall in *Eimeria maxima*: immunolight and electron microscopy. *Int J Parasitol* 33:1329–1340
- Gao, Y., Sun, P., Hu, D., Tang, X., Zhang, S., Shi, F., and Suo, X. (2024). Advancements in understanding chicken coccidiosis: from *Eimeria* biology to innovative control strategies. *One Health Advances*, 2(1), 1-19.
- Hassan, S. M., Zayeda, R., Elakany, H., Badr, S., Abou-Rawash, A., & Abd-Ellatieff, H. (2024). Anticoccidial activity of Aloe Vera L eafs' aqueous extract and vaccination against *Eimeria tenella*: pathological study in broilers. *Veterinary Research Communications*, 48(1), 403-416.
- Hegazi, A. G., Shanawany, E. E. E., El-Houssiny, A. S., Hassan, S. E., Desouky, H. M., El-Metenawy, T. M., & Abdel-Rahman, E. H. (2023). Attenuation of pathogenesis of *Eimeria stiedae* sporulated oocysts using Egyptian alginate propolis nanoparticles. *BMC veterinary research*, 19(1), 127.
- Jilo, S. A., Abadula, T. A., Abadura, S. Z., Gobana, R. H., Hasan, L. A., and Nair, S. P. (2022). Review on epidemiology, pathogenesis, treatment, control and prevention of gastrointestinal parasite of poultry.

- Kamal, E., Zaghawa, A., Salma, A., Nayel, M., and Dawoud, M. (2024). Risk Factors Associated with Common Infectious Diseases in Beef Cattle in Menofia Governorate. *Journal of Current Veterinary Research*, 6(1), 39-53.
- Kheirandish, R., Nourollahi-Fard, S.R., and Faryabi, Z. (2012). Prevalence and pathologic study of *Eimeria cameli* in slaughtered camels. *Eurasian J Vet Sci* 28:138–141
- Koudela, B., and Bokova, A. 1998. Coccidiosis in goats in the Czech Republic. *Veterinary Parasitology*. 76: 261–267
- Murshed, M., AL-Tamimi, J., Aljawdah, H., Mares, M. M., & Al-Quraishy, S. (2024). Effect of *Calotropis procera* extract against *Eimeria piriformis* oocyst-and sporozoite-infected rabbits. *Tropical Journal of Pharmaceutical Research*, 23(6).
- Safaa M. K. and May H. K. (2020) Traditional Diagnosis of *Eimeria* spp. in Fallow Deer at Middle Parts of Iraq. *The Iraqi Journal of Veterinary Medicine*, 44(E0), 94-99.
- Spencer, L., Bancroft, J., Bancroft, J., & Gamble, M. (2012). Tissue processing. *Bancroft's Theory and Practice of Histological Techniques*. 7th ed. Netherlands, Amsterdam: Elsevier Health Sciences, 105-23.
- Utebaeva, G., Berkinbay, O., and Tuganbay, A. (2021). Study of Prevalence and Associated Risk Factors of *Eimeria* sp., in Camels in Turkestan Region. *Archives of Razi Institute*, 76(5), 1419.
- Zainab, S. R., Shahzad, M. I., Mustafa, M. Z., Arshad, M., & Ruby, T. (2016). Prevalence of *Eimeria bovis* in cattles of Cholistan desert, Pakistan. *Journal of Biodiversity and Environmental Sciences*, 9, 94-98.